

Disclaimer: These policies are clinical editing payment decisions for Blue Cross of Blue Shield of Michigan and do not apply to Blue Care Network of Michigan.

Metabolite Markers of Thiopurines Testing

Policy Number: AHS – G2115 – Metabolite Markers of Thiopurines Testing	Prior Policy Name and Number, as applicable: <ul style="list-style-type: none"> AHS-G2115 Pharmacogenomic and Metabolite Markers for Thiopurines
Original Effective Date: 6/01/2022 Current Effective Date: 6/01/2022	

I. Policy Description

Thiopurines are a class of purine antimetabolite immunomodulators with diverse clinical applications in treatment of autoimmune disorders, transplant rejection, and acute lymphoblastic leukemia (Belmont, 2019; MacDermott, 2019). Their therapeutic efficacy, bone marrow toxicity, and liver toxicity have been reported to be related to levels of their downstream metabolites. Due to their complex metabolism, patient response varies considerably between individuals, both in achieving therapeutic drug levels as well as in developing adverse reactions (Bradford & Shih, 2011).

Please note that this policy discusses the monitoring of thiopurine metabolite levels in individuals. For guidance on pharmacogenetic testing prior to therapy, please refer to AHS-M2021 Pharmacogenetic Testing.

II. Related Policies

Policy Number	Policy Title
	Not Applicable

III. Indications and/or Limitations of Coverage

Application of coverage criteria is dependent upon an individual’s benefit coverage at the time of the request

- One-time phenotypic analysis of the enzyme TPMT **MEETS COVERAGE CRITERIA** in patients prior to initiating treatment with azathioprine (AZA), mercaptopurine (6-MP) or thioguanine (6-TG) OR in patients on thiopurine therapy

Disclaimer: These policies are clinical editing payment decisions for Blue Cross of Blue Shield of Michigan and do not apply to Blue Care Network of Michigan.

- with abnormal complete blood count (CBC) results that do not respond to dose reduction.
2. Monitoring of thiopurine metabolite levels in individuals with inflammatory bowel disease **MEETS COVERAGE CRITERIA** for the following indications:
 - a. To measure blood levels in individuals suspected of having toxic responses to AZA and/or 6-MP (e.g., hepatotoxicity or myelotoxicity);
 - b. To measure drug levels in individuals who have not responded to therapy (e.g., persistent fever, further weight loss, and bloody diarrhea).
 3. Monitoring of thiopurine metabolite levels in individuals with acute lymphoblastic leukemia **MEETS COVERAGE CRITERIA** in the following situations:
 - a. For patients showing signs of a lack of myelosuppression while on therapy
 - b. For patients with normal function for TPMT and NUDT15 who do not appear to tolerate thiopurines

The following does not meet coverage criteria due to a lack of available published scientific literature confirming that the test(s) is/are required and beneficial for the diagnosis and treatment of a patient's illness.

4. Phenotypic analysis of the enzyme TPMT **DOES NOT MEET COVERAGE CRITERIA** in all other situations.
5. Phenotypic analysis of the enzyme NUDT15 **DOES NOT MEET COVERAGE CRITERIA** in all other situations.
6. Analysis of the metabolite markers of azathioprine and 6-mercaptopurine, including 6-methyl-mercaptopurine ribonucleotides (6-MMRP) and 6-thioguanine nucleotides (6-TGN), **DOES NOT MEET COVERAGE CRITERIA** in all other situations.

IV. Scientific Background

The thiopurine drugs 6-mercaptopurine (6-MP), azathioprine (AZA), and thioguanine remain a mainstay of immunomodulator therapy (Belmont, 2019; Rubin, 2020; Tantisira, 2019). However, several metabolites of these drugs (particularly 6-thioguanine [6-TG] and 6-methylmercaptopurine [6-MMP]) have been associated with harmful side effects, such as lowered therapeutic efficacy, hepatotoxicity, bone marrow toxicity, and more. The management of these toxic metabolites is further complicated by the many polymorphisms (and therefore efficacy in metabolism) of the genes responsible for metabolizing these drugs.

Disclaimer: These policies are clinical editing payment decisions for Blue Cross of Blue Shield of Michigan and do not apply to Blue Care Network of Michigan.

Due to these toxic side effects, there has been significant investigation on monitoring of these metabolites to identify the optimal dose of a thiopurine for an individual patient. This process is called therapeutic drug monitoring (TDM) (MacDermott, 2019).

Two enzymes are responsible for catalyzing these reactions: thiopurine methyltransferase (TPMT) and hypoxanthine phosphoribosyl transferase. TPMT enzyme activity is a major factor determining AZA and 6-MP metabolism, and therefore 6-TG and 6-MMP levels. The majority of the population has wild type *TPMT* and normal enzyme activity, while 11% are heterozygous and have corresponding low TPMT enzyme activity, and 0.3% (1 in 300) have negligible activity (Lennard, Gibson, Nicole, & Lilleyman, 1993; Lennard, Van Loon, & Weinshilboum, 1989; Rubin, 2020). Intermediate and normal metabolizers can have up to a threefold difference in initial target doses of AZA and 6-MP to achieve therapeutic 6-TG concentrations (Gardiner, Geary, Begg, Zhang, & Barclay, 2008). Measurement of TPMT enzyme activity before instituting AZA or 6-MP may help prevent toxicity by identifying individuals with low or absent TPMT enzyme activity as well as identify those with higher than average TPMT activity who may remain refractory to conventional dosages (MacDermott, 2019). Dosing strategies involving such testing may be cost-effective (Cuffari, Dassopoulos, Turnbough, Thompson, & Bayless, 2004; Dubinsky et al., 2005; Winter et al., 2004). However, prediction of toxicity is not consistently reliable, as other enzymes are also likely to play a role in determining toxicity, such as glutathione-S-transferase (Stocco et al., 2007), and drug interactions must be taken into account (Dewit, Vanheuverzwyn, Desager, & Horsmans, 2002; Gilissen et al., 2005; Szumlanski & Weinshilboum, 1995). Thus, even though TPMT testing is recommended (Relling et al., 2019), a complete blood count (CBC), and also liver function tests, must still be obtained (Belmont, 2019; MacDermott, 2019).

Another enzyme that may contribute to thiopurine metabolism is nucleoside diphosphate-linked moiety X motif 15, NUDIX 15 (NUDT15). Variants in this enzyme's genotype and subsequent phenotype may lead to drastically reduced tolerance of 6-MP (Tantisira, 2019). Moriyama et al. proposed that NUDT15 variants cause thiopurines' mechanism of action to fail by preventing the thiopurine metabolites' incorporation into DNA. This causes these metabolites to remain active and therefore toxic (Moriyama et al., 2016). The frequency of these NUDT15 variants varies across populations, with the "poor metabolizer" phenotype reaching as high as 1 in 50 in East Asian populations. Despite the data indicating NUDT15's role in thiopurine toxicity, guidelines for its assessment have not reached a consensus, and expert opinions and practices are mixed (Tantisira, 2019).

Therapeutic drug monitoring (TDM) is the measurement of serum, plasma, or urinary concentrations of a given drug. This can be measured in a variety of ways, including high performance liquid chromatography (HPLC) or mass spectrometry approaches such as GC-MS (Eadie, 1998). TDM is proposed to allow a clinician to identify the "optimal" dose of a drug (such as a thiopurine) for a patient, thereby maximizing therapeutic efficacy and

Disclaimer: These policies are clinical editing payment decisions for Blue Cross of Blue Shield of Michigan and do not apply to Blue Care Network of Michigan.

minimizing harmful side effects. Non-TDM approaches typically involve starting at low doses, then adjusting if the patient is tolerating the drug well or poorly, whereas TDM takes a more proactive approach in managing dose (MacDermott, 2019). Several studies have attempted to identify standardized ranges of “optimal” metabolite concentrations. For example, the optimal concentration of the 6-TGN metabolite was found to be between 230 and 400 picomoles per 8×10^8 erythrocytes by Dubinsky et al. In that same study, bone marrow toxicity was found to correlate with levels above 400 picomoles per 8×10^8 erythrocytes (Dubinsky et al., 2000). Although there are potential limitations to TDM for thiopurines (such as intra-individual variability, lack of correlation with toxicity for 6-MMP, and so on), TDM used in conjunction with TPMT and NUDT15 assessment may allow clinicians to increase the therapeutic efficacy of thiopurines (MacDermott, 2019; Tantisira, 2019).

Clinical Validity and Utility

Haines et al. (2011) performed a retrospective study of the utility of measuring thiopurine metabolites in “inadequately controlled” inflammatory bowel disease (IBD). 63 patients with IBD were included, and these patients were treated with AZA or 6-mercaptopurine. On weight-based criteria, 50% patients were underdosed. However, metabolite study suggested that “7 (11%) patients were noncompliant, 18 (29%) were being underdosed, 33 (52%) were refractory to treatment with either appropriate (41%) or elevated (11%) metabolite concentrations, and 6 (10%) had a raised 6-methyl mercaptopurine: 6-thioguanine nucleotide ratio consistent with aberrant thiopurine metabolism”. The clinical outcome of 87% of patients improved when the treatment was shifted to a metabolite-based algorithm, whereas 3 of 17 patients improved when the discordant action was taken. The authors concluded that “Thiopurine metabolite testing is a potentially powerful tool for optimizing thiopurine usage in IBD” (Haines et al., 2011).

Lee et al. (2017) evaluated 165 patients undergoing thiopurine treatment for Crohn’s Disease. Thiopurine metabolite levels were measured, and both *TPMT* and *NUDT15* were genotyped. The authors found 95 patients responded to treatment whereas 45 did not. The median 6-TGN (the primary metabolite of 6-thioguanine) was significantly higher in responders than nonresponders. At a 6-TGN level of 230 pmol/ 8×10^8 blood cells, the odds ratio was 4.63 for responders to nonresponders. *NUDT15* variants were also found to be associated with “severe, early, leukopenia” with an average reduction of 88.2% from baseline white blood cell count at 4 weeks. The authors concluded that their findings “support the role of therapeutic drug monitoring in thiopurine maintenance treatment to optimize thiopurine therapy, especially, for non-responding CD patients” (Lee et al., 2017).

Spencer, Norris, Williams, and Dubinsky (2019) compared “standard” and “optimized” thiopurine dosing regimens in 216 pediatric IBD patients. The “optimal” level was decided at “6-TGN >235 pmol/ 8×10^8 RBC”, and the metabolite levels were correlated between the primary outcome of “steroid-free clinical remission (SFR)”. Both groups were found to have

Disclaimer: These policies are clinical editing payment decisions for Blue Cross of Blue Shield of Michigan and do not apply to Blue Care Network of Michigan.

similar initial and 6-month metabolite levels. SFR was achieved in 74% of the 180 patients on thiopurines at 6 months. The authors concluded that “steroid-free clinical remission and 6-TGN levels at 6 months were no different between a standardized, fixed dosing strategy and a metabolite-driven, optimized dosing strategy” (Spencer et al., 2019).

Meijer et al. (2017) evaluated the effects of thiopurine metabolites on clinical signs and if patient characteristics affected metabolite generation. 940 “laboratory findings” from 424 patients were examined. 6-TGN (a metabolite of azathioprine [AZA] and mercaptopurine) was found to negatively correlate with RBC count, WBC count, and neutrophil count. However, in patients using 6-thioguanine, those 6-TGN concentrations correlated positively with WBC count. An inverse correlation was observed between age and 6-TGN concentrations in AZA or 6-thioguanine patients, as well as between body mass index and 6-TGN in AZA or mercaptopurine patients. The authors concluded that “thiopurine derivative therapy influenced bone marrow production and the size of red blood cells. Age and body mass index were important pharmacokinetic factors in the generation of 6-TGN” (Meijer et al., 2017).

Estevinho et al. (2017) performed a meta-analysis to “assess the clinical value of 6-thioguanine nucleotide thresholds; and ii) to compare mean 6-thioguanine nucleotide concentrations between patients in clinical remission vs. those with active disease.” A total of 22 records were used in cut-off comparisons and 12 were used in the 6-thioguanine nucleotide mean differences analysis. The authors calculated the global odds ratio for remission in patients with 6-thioguanine nucleotides above predefined thresholds to be 3.95. The authors also found an odds ratio for remission of 2.25 for the $235 \text{ pmol}/8 \times 10^8 \text{ RBC}$ threshold, and an odds ratio of 4.71 for the $250 \text{ pmol}/8 \times 10^8 \text{ RBC}$ threshold. Finally, the authors found a “pooled difference” of $63.37 \text{ pmol}/8 \times 10^8 \text{ RBC}$ between patients in clinical remission and those not in remission. Overall, the authors concluded that the study reinforced the link between 6-thioguanine nucleotide levels and clinical remission in inflammatory bowel diseases (Estevinho et al., 2017).

Toksvang et al. (2019) performed a meta-analysis focusing on “incidence of hepatotoxicity in patients [with childhood acute lymphoblastic leukaemia, ALL or inflammatory bowel disease, IBD] treated with 6TG [6-thioguanine]”. 42 reports were included, further broken down into “four randomised controlled trials (RCTs) including 3,993 patients, 20 observational studies including 796 patients, and 18 case reports including 60 patients”. The authors measured hepatotoxicity by “sinusoidal obstruction syndrome”, which occurred in 9-25% of ALL patients in two of the four RCTs at a dosage of 40–60 $\text{mg}/\text{m}^2/\text{day}$. The authors also noted that at a dosage of 23 $\text{mg}/\text{m}^2/\text{day}$, nodular regenerative hyperplasia (NRH) occurred in 14% of IBD patients. At a dosage of 12 $\text{mg}/\text{m}^2/\text{day}$, NRH occurred in 6% of IBD patients, which was noted to be similar to background incidence. The authors therefore concluded that doses at or under 12 $\text{mg}/\text{m}^2/\text{day}$ can “probably be considered safe” (Toksvang et al., 2019).

Disclaimer: These policies are clinical editing payment decisions for Blue Cross of Blue Shield of Michigan and do not apply to Blue Care Network of Michigan.

Zhu et al. (2019) evaluated the “predictive sensitivity based on 6TGN [6-thioguanine nucleotide] by subgrouping patients according to their *NUDT15* R139C genotypes”. The authors included 411 patients with Crohn’s Disease. Two subgroups of *NUDT15* genotypes were created, “CC” (n = 342) and “CT” (n = 65), with the final four patients harboring a TT genotype. Thiopurine-induced leukopenia (TIL) was the primary clinical endpoint measured. The authors found that of the 342 patients with a CC genotype, only 35 developed TIL (10.2%), but of the 65 CT patients, 33 developed TIL (50.2%). All four of the TT patients developed TIL. The authors also found that in both CC and CT genotypes, the median 6TGN level was higher in patients with TIL than patients without TIL (for CC, 474.8 pmol/8 × 10⁸ RBC vs 306.0 pmol/8 × 10⁸ RBC, for CT 291.7 / 8 × 10⁸ RBC vs 217.6/8 × 10⁸ RBC). From this data, the authors calculated the “cut-off” (a threshold to identify an optimal amount of cases) of the CT genotype to be 319.2 pmol/8 × 10⁸ RBC and the cut-off for CC to be 411.5 pmol/8 × 10⁸ RBC). Overall, the authors concluded that “The predictive sensitivity of TIL based on 6TGN is dramatically increased after subgrouping according to *NUDT15* R139C genotypes. Applying 6TGN cut-off levels to adjust thiopurine therapies based on *NUDT15* is strongly recommended” (Zhu et al., 2019).

V. Guidelines and Recommendations

National Comprehensive Cancer Network (NCCN, 2019, 2020, 2021a, 2021b)

In version 2 of the 2021 guidelines for Pediatric Acute Lymphoblastic Leukemia, the NCCN recommends that “for patients homozygous for normal function TPMT and *NUDT15*, who do not appear to tolerate thiopurines, consider measuring erythrocyte thiopurine metabolites and/or erythrocyte TPMT activity. Genetic testing may fail to identify rare or previously undiscovered no function alleles.” The NCCN also writes that “genetic testing for no function alleles of TPMT and *NUDT15* should be considered prior to the initiation of thiopurine therapy” (NCCN, 2019, 2021b)

In version 2 of the 2021 guidelines for Acute Lymphoblastic Leukemia, the NCCN notes that “quantification of 6-MP metabolites can be very useful in determining whether lack of myelosuppression is due to non-compliance or hypermetabolism”

The 2021 guidelines also state, “for patients receiving 6-MP, consider testing for TPMT gene polymorphisms, particularly in patients who develop severe neutropenia after starting 6-MP. Testing for both TPMT and *NUDT15* variant status should be considered, especially for patients of East Asian origin” (NCCN, 2021a).

Toronto Ulcerative Colitis Consensus Group/American College of Gastroenterology (ACG) (Bressler et al., 2015)

Bressler et al. (2015) published clinical practice guidelines for the medical management of non-hospitalized ulcerative colitis on behalf of the Toronto Ulcerative Colitis Consensus

Disclaimer: These policies are clinical editing payment decisions for Blue Cross of Blue Shield of Michigan and do not apply to Blue Care Network of Michigan.

Group, which reaffirmed recommendations from the American College of Gastroenterology, Practice Parameters Committee (Kornbluth & Sachar, 2010) for thiopurine therapy (Bressler et al., 2015). The authors stated that "...a TPMT assay is necessary before initiation of treatment to identify patients at risk for severe dose-dependent myelosuppression...therefore, thiopurine metabolite levels may be helpful to guide therapy. Note that TPMT testing does not replace the need for mandatory monitoring of complete blood cell count" (Bressler et al., 2015).

American College of Gastroenterology (ACG) (Lichtenstein et al., 2018; Rubin, Ananthakrishnan, Siegel, Sauer, & Long, 2019)

The ACG published a guideline for the "Management of Crohn's Disease in Adults". Their relevant recommendations include:

"Thiopurine methyltransferase (TPMT) testing should be considered before initial use of azathioprine or 6-mercaptopurine to treat patients with Crohn's disease (strong recommendation, low level of evidence)" (Lichtenstein et al., 2018).

The ACG also published a guideline for ulcerative colitis (UC) in adults. Their relevant recommendations include:

"The patient with nonresponse or loss of response to therapy should be assessed with therapeutic drug monitoring to identify the reason for lack of response and whether to optimize the existing therapy or to select an alternate therapy."

"There is insufficient evidence supporting a benefit for proactive therapeutic drug monitoring in all unselected patients with UC in remission" (Rubin et al., 2019).

North American Society for Pediatric Gastroenterology, Hepatology and Nutrition (NASPGHAN) Committee (Benkov et al., 2013)

In 2013, the North American Society for Pediatric Gastroenterology, Hepatology and Nutrition (NASPGHAN) Committee on Inflammatory Bowel Disease published consensus recommendations on the role of TPMT and thiopurine metabolite testing in pediatric IBD (Benkov et al., 2013) The recommendations included the following:

- "TPMT testing is recommended before initiation of TPs to identify individuals who are homozygote recessive or have extremely low TPMT activity, with the latter having more reliability than the former. (HIGH)."
- "Individuals who are homozygous recessive or have extremely low TPMT activity should avoid use of TPs because of concerns for significant leukopenia. (HIGH)"

Disclaimer: These policies are clinical editing payment decisions for Blue Cross of Blue Shield of Michigan and do not apply to Blue Care Network of Michigan.

- “TPMT testing does not predict all cases of leukopenia and has no value to predict hypersensitivity adverse effects such as pancreatitis. Any potential value to reduce the risk of malignancy has not been studied. All individuals on TPs should have routine monitoring with CBC and WBC count differential to evaluate for leukopenia regardless of TPMT testing results. (HIGH)”
- “Metabolite testing can be used to determine adherence to TP therapy. (HIGH)”
- “Metabolite testing can be used to guide dose increases or modifications in patients with active disease. Consideration would include either increasing the dose, changing therapy or for those with elevated transaminases or an elevated 6-MMP, using adjunctive allopurinol to help raise 6-TG metabolites and suppress formation of 6-MMP. (MODERATE)”
- “Routine and repetitive metabolite testing has little or no role in patients who are doing well and taking an acceptable dose of a TP. (MODERATE)”

American Gastroenterological Association (AGA) (Joseph D. Feuerstein et al., 2020; J. D. Feuerstein, Nguyen, Kupfer, Falck-Ytter, & Singh, 2017)

In 2017, the AGA published guidelines (J. D. Feuerstein et al., 2017) on Therapeutic Drug Monitoring in Inflammatory Bowel Disease which recommend:

- “In adult patients treated with thiopurines with active IBD or adverse effects thought to be due to thiopurine toxicity, the AGA suggests reactive thiopurine metabolite monitoring to guide treatment changes.”
- “In adult patients with quiescent IBD treated with thiopurines, the AGA suggests against routine thiopurine metabolite monitoring.”

The AGA published an Institute Technical Review on the Role of Therapeutic Drug Monitoring in the Management of Inflammatory Bowel Diseases in the same year. In it, they note that IBD patients treated with thiopurines may benefit from reactive TDM to guide treatment changes (Vande Casteele, Herfarth, Katz, Falck-Ytter, & Singh, 2017).

In the 2020 AGA guidelines for “Management of Moderate to Severe Ulcerative Colitis”, the AGA remarks that “therapeutic drug monitoring to guide the use of biologic therapy has been addressed in separate AGA guidelines”. The “separate AGA guidelines” refer to the 2017 edition above (Joseph D. Feuerstein et al., 2020).

Clinical Pharmacogenetics Implementation Consortium (CPIC, 2020; Relling et al., 2019)

In their guideline for “Thiopurine Dosing Based on TPMT and NUDT15 Genotypes”, CPIC

Disclaimer: These policies are clinical editing payment decisions for Blue Cross of Blue Shield of Michigan and do not apply to Blue Care Network of Michigan.

notes that “mercaptopurine and azathioprine are generally used for nonmalignant immunologic disorders, mercaptopurine for lymphoid malignancies, and thioguanine for myeloid leukemias”. However, CPIC also writes that “variants in NUDT15 have been identified that strongly influence thiopurine tolerance in patients with acute lymphoblastic leukemia (ALL) and those with inflammatory bowel diseases” (Relling et al., 2019).

Although CPIC published an official guideline regarding TPMT and NUDT15 testing in 2019, an update was posted to their website in April 2020. That update is as follows:

- “For TPMT and NUDT15 indeterminate phenotypes, (i.e. combination of uncertain and/or unknown function alleles):
 - TPMT indeterminate: Consider evaluating TPMT erythrocyte activity to assess TPMT phenotype.
 - NUDT15 indeterminate: If thiopurines are required and NUDT15 status is unknown, monitor closely for toxicity” (CPIC, 2020).

British Society of Gastroenterology (BSG) (Lamb et al., 2019)

The BSG published “consensus guidelines” on management of inflammatory bowel disease in adults. They recommend checking TPMT status in “all patients considered for thiopurine therapy”. They also recommend testing the NUDT15 genotype if “available”.

The BSG also writes that thiopurine metabolites should be checked if a patient experiences myelotoxicity as a side effect. Similarly, if a patient demonstrates “newly abnormal LFTs [liver function tests]”, thiopurine metabolites should be checked.

Overall, the BSG writes that thiopurine metabolites can be used to “optimize drug dosing” and “suggest that metabolite monitoring may be used for those with inadequate response to therapy or toxicity, but should not be a substitute for routine monitoring blood tests” (Lamb et al., 2019).

Canadian Association of Gastroenterology (Mack et al., 2019)

The Canadian Association of Gastroenterology published a guideline on “the Medical Management of Pediatric Luminal Crohn’s Disease.” The guidelines “suggested that testing for TPMT by genotype or enzymatic activity be done prior to initiating thiopurine therapy to guide dosing” (Mack et al., 2019).

An additional guideline for [Adult] Luminal Crohn’s Disease specifies “because some patients may have low or absent levels of the enzyme (thiopurine methyltransferase (TPMT) needed to

Disclaimer: These policies are clinical editing payment decisions for Blue Cross of Blue Shield of Michigan and do not apply to Blue Care Network of Michigan.

metabolize thiopurines, a TPMT assay should be performed before initiation of treatment to identify patients at risk for severe toxicity” (Panaccione et al., 2018)

National Institute for Health and Care Excellence (NICE, 2019)

NICE released guidelines on Crohn’s Disease in 2019. In it, they recommend to “Monitor the effects of azathioprine, mercaptopurine, and methotrexate as advised in the British national formulary (BNF) or British national formulary for children (BNFC)” (NICE, 2019).

European Crohn's and Colitis Organization and European Society of Paediatric Gastroenterology, Hepatology and Nutrition (Turner et al., 2018)

These joint guidelines note that “measuring...6-TG and 6-MMP levels after 2–3 months, may aid in optimizing thiopurine dosing”. Measuring thiopurine metabolites is recommended in the following scenarios:

- In patients with incomplete response on stable thiopurine dosage
- In patients who present with leucopenia or elevated transaminases
- When poor compliance is suspected (Turner et al., 2018).

VI. State and Federal Regulations, as applicable

DISCLAIMER: If there is a conflict between this Policy and any relevant, applicable government policy for a particular member [e.g. Local Coverage Determinations (LCDs) or National Coverage Determinations (NCDs) for Medicare and/or state coverage for Medicaid], then the government policy will be used to make the determination. For the most up-to-date Medicare policies and coverage, please visit the Medicare search website <http://www.cms.gov/medicare-coverage-database/overview-and-quick-search.aspx>. For the most up-to-date Medicaid policies and coverage, please visit the applicable state Medicaid website.

Many labs have developed specific tests that they must validate and perform in house. These laboratory-developed tests (LDTs) are regulated by the Centers for Medicare and Medicaid (CMS) as high-complexity tests under the Clinical Laboratory Improvement Amendments of 1988 (CLIA '88). FDA clearance or approval is not currently required for clinical use.

LDTs may be covered when Medicare coverage criteria are met. Refer to the [Medicare Claims Processing Manual Chapter 12 - Physicians/Nonphysician Practitioners](#) for details. In addition, Local Coverage Determinations (LCDs)/Local Coverage Articles (LCAs) are available for

Disclaimer: These policies are clinical editing payment decisions for Blue Cross of Blue Shield of Michigan and do not apply to Blue Care Network of Michigan.

reference. Compliance with these policies is mandatory where applicable. Find these LCDs/LCAs at <https://www.cms.gov/medicare-coverage-database/new-search/search.aspx> (CMS, 2021).

There is no national coverage determination for this policy. When a national coverage determination does not exist, coverage decisions are left to local Medicare carrier moderation and judgment.

VII. Applicable CPT/HCPCS Procedure Codes

Code Number	Code Description
80299	Quantitation of therapeutic drug, not elsewhere specified
82657	Enzyme activity in blood cells, cultured cells, or tissue, not elsewhere specified; nonradioactive substrate, each specimen

Current Procedural Terminology© American Medical Association. All Rights reserved.

Procedure codes appearing in Medical Policy documents are included only as a general reference tool for each policy. They may not be all-inclusive.

VIII. Evidence-based Scientific References

- Belmont, H. M. (2019). Pharmacology and side effects of azathioprine when used in rheumatic diseases - UpToDate. In P. L. Romain (Ed.), *UpToDate*. Retrieved from https://www.uptodate.com/contents/pharmacology-and-side-effects-of-azathioprine-when-used-in-rheumatic-diseases?source=see_link§ionName=Pharmacogenetics%20and%20TPMT%20testing&anchor=H4#H4
- Benkov, K., Lu, Y., Patel, A., Rahhal, R., Russell, G., & Teitelbaum, J. (2013). Role of thiopurine metabolite testing and thiopurine methyltransferase determination in pediatric IBD. *J Pediatr Gastroenterol Nutr*, *56*(3), 333-340. doi:10.1097/MPG.0b013e3182844705
- Bradford, K., & Shih, D. Q. (2011). Optimizing 6-mercaptopurine and azathioprine therapy in the management of inflammatory bowel disease. *World J Gastroenterol*, *17*(37), 4166-4173. doi:10.3748/wjg.v17.i37.4166
- Bressler, B., Marshall, J. K., Bernstein, C. N., Bitton, A., Jones, J., Leontiadis, G. I., . . . Feagan, B. (2015). Clinical practice guidelines for the medical management of nonhospitalized

Disclaimer: These policies are clinical editing payment decisions for Blue Cross of Blue Shield of Michigan and do not apply to Blue Care Network of Michigan.

- ulcerative colitis: the Toronto consensus. *Gastroenterology*, 148(5), 1035-1058.e1033. doi:10.1053/j.gastro.2015.03.001
- CMS. (2021). Medicare Coverage Database. Retrieved from <https://www.cms.gov/medicare-coverage-database/new-search/search.aspx>. <https://www.cms.gov/medicare-coverage-database/new-search/search.aspx>
- CPIC. (2020). CPIC® Guideline for Thiopurines and TPMT and NUDT15. Retrieved from <https://cpicpgx.org/guidelines/guideline-for-thiopurines-and-tpmt/>
- Cuffari, C., Dassopoulos, T., Turnbough, L., Thompson, R. E., & Bayless, T. M. (2004). Thiopurine methyltransferase activity influences clinical response to azathioprine in inflammatory bowel disease. *Clin Gastroenterol Hepatol*, 2(5), 410-417. Retrieved from <http://dx.doi.org/>
- Dewit, O., Vanheuverzwyn, R., Desager, J. P., & Horsmans, Y. (2002). Interaction between azathioprine and aminosalicylates: an in vivo study in patients with Crohn's disease. *Aliment Pharmacol Ther*, 16(1), 79-85. Retrieved from <http://dx.doi.org/>
- Dubinsky, M. C., Lamothe, S., Yang, H. Y., Targan, S. R., Sinnett, D., Theoret, Y., & Seidman, E. G. (2000). Pharmacogenomics and metabolite measurement for 6-mercaptopurine therapy in inflammatory bowel disease. *Gastroenterology*, 118(4), 705-713. Retrieved from <http://dx.doi.org/>
- Dubinsky, M. C., Reyes, E., Ofman, J., Chiou, C. F., Wade, S., & Sandborn, W. J. (2005). A cost-effectiveness analysis of alternative disease management strategies in patients with Crohn's disease treated with azathioprine or 6-mercaptopurine. *Am J Gastroenterol*, 100(10), 2239-2247. doi:10.1111/j.1572-0241.2005.41900.x
- Eadie, M. J. (1998). Therapeutic drug monitoring--antiepileptic drugs. *Br J Clin Pharmacol*, 46(3), 185-193. doi:10.1046/j.1365-2125.1998.00769.x
- Estevinho, M. M., Afonso, J., Rosa, I., Lago, P., Trindade, E., Correia, L., . . . Magro, F. (2017). A Systematic Review and Meta-Analysis of 6-Thioguanine Nucleotide Levels and Clinical Remission in Inflammatory Bowel Disease. *J Crohns Colitis*, 11(11), 1381-1392. doi:10.1093/ecco-jcc/jjx089
- Feuerstein, J. D., Isaacs, K. L., Schneider, Y., Siddique, S. M., Falck-Ytter, Y., Singh, S., . . . Siddique, S. M. (2020). AGA Clinical Practice Guidelines on the Management of Moderate to Severe Ulcerative Colitis. *Gastroenterology*, 158(5), 1450-1461. doi:10.1053/j.gastro.2020.01.006
- Feuerstein, J. D., Nguyen, G. C., Kupfer, S. S., Falck-Ytter, Y., & Singh, S. (2017). American Gastroenterological Association Institute Guideline on Therapeutic Drug Monitoring in Inflammatory Bowel Disease. *Gastroenterology*, 153(3), 827-834. doi:10.1053/j.gastro.2017.07.032
- Gardiner, S. J., Gearry, R. B., Begg, E. J., Zhang, M., & Barclay, M. L. (2008). Thiopurine dose in intermediate and normal metabolizers of thiopurine methyltransferase may differ three-fold. *Clin Gastroenterol Hepatol*, 6(6), 654-660; quiz 604. doi:10.1016/j.cgh.2008.02.032

Disclaimer: These policies are clinical editing payment decisions for Blue Cross of Blue Shield of Michigan and do not apply to Blue Care Network of Michigan.

- Gilissen, L. P., Bierau, J., Derijks, L. J., Bos, L. P., Hooymans, P. M., van Gennip, A., . . . Engels, L. G. (2005). The pharmacokinetic effect of discontinuation of mesalazine on mercaptopurine metabolite levels in inflammatory bowel disease patients. *Aliment Pharmacol Ther*, 22(7), 605-611. doi:10.1111/j.1365-2036.2005.02630.x
- Haines, M. L., Ajlouni, Y., Irving, P. M., Sparrow, M. P., Rose, R., Geary, R. B., & Gibson, P. R. (2011). Clinical usefulness of therapeutic drug monitoring of thiopurines in patients with inadequately controlled inflammatory bowel disease. *Inflamm Bowel Dis*, 17(6), 1301-1307. doi:10.1002/ibd.21458
- Kornbluth, A., & Sachar, D. B. (2010). Ulcerative colitis practice guidelines in adults: American College Of Gastroenterology, Practice Parameters Committee. *Am J Gastroenterol*, 105(3), 501-523; quiz 524. doi:10.1038/ajg.2009.727
- Lamb, C. A., Kennedy, N. A., Raine, T., Hendy, P. A., Smith, P. J., Limdi, J. K., . . . Hawthorne, A. B. (2019). British Society of Gastroenterology consensus guidelines on the management of inflammatory bowel disease in adults. *Gut*, 68(Suppl 3), s1-s106. doi:10.1136/gutjnl-2019-318484
- Lee, J. H., Kim, T. J., Kim, E. R., Hong, S. N., Chang, D. K., Choi, L. H., . . . Kim, Y. H. (2017). Measurements of 6-thioguanine nucleotide levels with TPMT and NUDT15 genotyping in patients with Crohn's disease. *PLoS One*, 12(12), e0188925. doi:10.1371/journal.pone.0188925
- Lennard, L., Gibson, B. E., Nicole, T., & Lilleyman, J. S. (1993). Congenital thiopurine methyltransferase deficiency and 6-mercaptopurine toxicity during treatment for acute lymphoblastic leukaemia. *Arch Dis Child*, 69(5), 577-579. Retrieved from <http://dx.doi.org/>
- Lennard, L., Van Loon, J. A., & Weinshilboum, R. M. (1989). Pharmacogenetics of acute azathioprine toxicity: relationship to thiopurine methyltransferase genetic polymorphism. *Clin Pharmacol Ther*, 46(2), 149-154. Retrieved from <http://dx.doi.org/>
- Lichtenstein, G. R., Loftus, E. V., Isaacs, K. L., Regueiro, M. D., Gerson, L. B., & Sands, B. E. (2018). ACG Clinical Guideline: Management of Crohn's Disease in Adults. *Official journal of the American College of Gastroenterology | ACG*, 113(4). Retrieved from https://journals.lww.com/ajg/Fulltext/2018/04000/ACG_Clinical_Guideline_Management_of_Crohn_s.10.aspx
- MacDermott, R. (2019). 6-mercaptopurine (6-MP) metabolite monitoring and TPMT testing in the treatment of inflammatory bowel disease with 6-MP or azathioprine - UpToDate. In K. Robson (Ed.), *UpToDate*. Retrieved from https://www.uptodate.com/contents/6-mercaptopurine-6-mp-metabolite-monitoring-and-tpmt-testing-in-the-treatment-of-inflammatory-bowel-disease-with-6-mp-or-azathioprine?source=see_link§ionName=TPMT%20genotyping&anchor=H3#H3
- Mack, D. R., Benchimol, E. I., Critch, J., deBruyn, J., Tse, F., Moayyedi, P., . . . Griffiths, A. (2019). Canadian Association of Gastroenterology Clinical Practice Guideline for the

Disclaimer: These policies are clinical editing payment decisions for Blue Cross of Blue Shield of Michigan and do not apply to Blue Care Network of Michigan.

- Medical Management of Pediatric Luminal Crohn's Disease. *J Can Assoc Gastroenterol*, 2(3), e35-e63. doi:10.1093/jcag/gwz018
- Meijer, B., Wilhelm, A. J., Mulder, C. J. J., Bouma, G., van Bodegraven, A. A., & de Boer, N. K. H. (2017). Pharmacology of Thiopurine Therapy in Inflammatory Bowel Disease and Complete Blood Cell Count Outcomes: A 5-Year Database Study. *Ther Drug Monit*, 39(4), 399-405. doi:10.1097/ftd.0000000000000414
- Moriyama, T., Nishii, R., Perez-Andreu, V., Yang, W., Klusmann, F. A., Zhao, X., . . . Yang, J. J. (2016). NUDT15 polymorphisms alter thiopurine metabolism and hematopoietic toxicity. *Nat Genet*, 48(4), 367-373. doi:10.1038/ng.3508
- NCCN. (2019). Pediatric Acute Lymphoblastic Leukemia, Version 2. Retrieved from https://www.nccn.org/professionals/physician_gls/pdf/ped_all.pdf
- NCCN. (2020). Acute Lymphoblastic Leukemia. Retrieved from https://www.nccn.org/professionals/physician_gls/pdf/all.pdf
- NCCN. (2021a). Acute Lymphoblastic Leukemia. Retrieved from https://www.nccn.org/professionals/physician_gls/pdf/all.pdf
- NCCN. (2021b). Pediatric Acute Lymphoblastic Leukemia, Version 2. Retrieved from https://www.nccn.org/professionals/physician_gls/pdf/ped_all.pdf
- NICE. (2019). Crohn's disease: management. Retrieved from <https://www.nice.org.uk/guidance/ng129/chapter/Recommendations>
- Panaccione, R., Steinhart, A. H., Bressler, B., Khanna, R., Marshall, J. K., Targownik, L., . . . Bernstein, C. N. (2018). Canadian Association of Gastroenterology Clinical Practice Guideline for the Management of Luminal Crohn's Disease. *Journal of the Canadian Association of Gastroenterology*, 2(3), e1-e34. doi:10.1093/jcag/gwz019
- Relling, M. V., Schwab, M., Whirl-Carrillo, M., Suarez-Kurtz, G., Pui, C. H., Stein, C. M., . . . Yang, J. J. (2019). Clinical Pharmacogenetics Implementation Consortium Guideline for Thiopurine Dosing Based on TPMT and NUDT15 Genotypes: 2018 Update. *Clin Pharmacol Ther*, 105(5), 1095-1105. doi:10.1002/cpt.1304
- Rubin, D. T. (2020). Thiopurines: Pretreatment testing and approach to therapeutic drug monitoring for adults with inflammatory bowel disease. Retrieved from https://www.uptodate.com/contents/thiopurines-pretreatment-testing-and-approach-to-therapeutic-drug-monitoring-for-adults-with-inflammatory-bowel-disease?search=thiopurines&source=search_result&selectedTitle=2~44&usage_type=defaul&display_rank=2
- Rubin, D. T., Ananthkrishnan, A. N., Siegel, C. A., Sauer, B. G., & Long, M. D. (2019). ACG Clinical Guideline: Ulcerative Colitis in Adults. *Official journal of the American College of Gastroenterology | ACG*, 114(3). Retrieved from https://journals.lww.com/ajg/Fulltext/2019/03000/ACG_Clinical_Guideline_Ulcerative_Colitis_in.10.aspx

Disclaimer: These policies are clinical editing payment decisions for Blue Cross of Blue Shield of Michigan and do not apply to Blue Care Network of Michigan.

- Spencer, E., Norris, E., Williams, C., & Dubinsky, M. C. (2019). The Impact of Thiopurine Metabolite Monitoring on the Durability of Thiopurine Monotherapy in Pediatric IBD. *Inflamm Bowel Dis*, 25(1), 142-149. doi:10.1093/ibd/izy216
- Stocco, G., Martellosi, S., Barabino, A., Decorti, G., Bartoli, F., Montico, M., . . . Ventura, A. (2007). Glutathione-S-transferase genotypes and the adverse effects of azathioprine in young patients with inflammatory bowel disease. *Inflamm Bowel Dis*, 13(1), 57-64. doi:10.1002/ibd.20004
- Szumanski, C. L., & Weinshilboum, R. M. (1995). Sulphasalazine inhibition of thiopurine methyltransferase: possible mechanism for interaction with 6-mercaptopurine and azathioprine. *Br J Clin Pharmacol*, 39(4), 456-459. Retrieved from <http://dx.doi.org/>
- Tantisira, K., Weiss, Scott. (2019). Overview of pharmacogenomics. Retrieved from https://www.uptodate.com/contents/overview-of-pharmacogenomics?sectionName=Thiopurines%20and%20polymorphisms%20in%20TPMT%20and%20NUDT15&search=thioguanine&topicRef=4524&anchor=H6&source=se_e_link#H6
- Toksvang, L. N., Schmidt, M. S., Arup, S., Larsen, R. H., Frandsen, T. L., Schmiegelow, K., & Rank, C. U. (2019). Hepatotoxicity during 6-thioguanine treatment in inflammatory bowel disease and childhood acute lymphoblastic leukaemia: A systematic review. *PLoS One*, 14(5), e0212157. doi:10.1371/journal.pone.0212157
- Turner, D., Ruemmele, F. M., Orlanski-Meyer, E., Griffiths, A. M., de Carpi, J. M., Bronsky, J., . . . Russell, R. K. (2018). Management of Paediatric Ulcerative Colitis, Part 1: Ambulatory Care-An Evidence-based Guideline From European Crohn's and Colitis Organization and European Society of Paediatric Gastroenterology, Hepatology and Nutrition. *J Pediatr Gastroenterol Nutr*, 67(2), 257-291. doi:10.1097/mpg.0000000000002035
- Vande Casteele, N., Herfarth, H., Katz, J., Falck-Ytter, Y., & Singh, S. (2017). American Gastroenterological Association Institute Technical Review on the Role of Therapeutic Drug Monitoring in the Management of Inflammatory Bowel Diseases. *Gastroenterology*, 153(3), 835-857.e836. doi:10.1053/j.gastro.2017.07.031
- Winter, J., Walker, A., Shapiro, D., Gaffney, D., Spooner, R. J., & Mills, P. R. (2004). Cost-effectiveness of thiopurine methyltransferase genotype screening in patients about to commence azathioprine therapy for treatment of inflammatory bowel disease. *Aliment Pharmacol Ther*, 20(6), 593-599. doi:10.1111/j.1365-2036.2004.02124.x
- Zhu, X., Chao, K., Li, M., Xie, W., Zheng, H., Zhang, J. X., . . . Wang, X. D. (2019). Nucleoside diphosphate-linked moiety X-type motif 15 R139C genotypes impact 6-thioguanine nucleotide cut-off levels to predict thiopurine-induced leukopenia in Crohn's disease patients. *World J Gastroenterol*, 25(38), 5850-5861. doi:10.3748/wjg.v25.i38.5850

IX. Revision History



Nonprofit corporations and independent licensees
of the Blue Cross and Blue Shield Association

Disclaimer: These policies are clinical editing payment decisions for Blue Cross of Blue Shield of Michigan and do not apply to Blue Care Network of Michigan.

Effective Date	Summary
06/01/2022	Initial Policy Implementation