

Allergen Testing

Policy Number: AHS – G2031 – Allergen Testing	Prior Policy Name and Number, as applicable:
Policy Revision Date: 08/01/2024 Initial Policy Effective date 11/04/2021	

POLICY DESCRIPTION | RELATED POLICIES | INDICATIONS AND/OR LIMITATIONS OF COVERAGE | TABLE OF TERMINOLOGY | SCIENTIFIC BACKGROUND | GUIDELINES AND RECOMMENDATIONS | APPLICABLE STATE AND FEDERAL REGULATIONS | APPLICABLE CPT/HCPCS PROCEDURE CODES | EVIDENCE-BASED SCIENTIFIC REFERENCES | REVISION HISTORY

I. Policy Description

Allergic disease is characterized by inappropriate or exaggerated rated immune reactions to foreign antigens (allergens) that are generally innocuous to most people, but when introduced into a genetically-predisposed individual, elicit a hypersensitivity reaction (Hamilton, 2023). Hypersensitivity reactions can be classified into four types, two of which are associated with allergy, type I immediate immunoglobulin E (IgE) reactions and type IV T cell mediated reactions (Chang & Guarderas, 2018). Type I reactions involve the formation of IgE antibodies specific to the allergen. When the subject is re-exposed to that allergen, the allergen binds multiple IgE molecules, resulting in the release of an array of inflammatory mediators, including histamines, that precipitate the symptoms of allergic disease (Hamilton, 2023).

Allergen testing in serum is designed to detect the presence of allergen specific IgE. A positive test for allergen specific IgE confirms the presence of the antibody only. Actual reactivity must be determined by history or supervised challenge (Kowal & DuBuske, 2022). Several diagnostic procedures have been developed to elicit and assess hypersensitivity reactions including epicutaneous, intradermal, patch, bronchial, exercise, and ingestion challenge tests (Bernstein et al., 2008).

II. Related Policies

Policy Number	Policy Title
AHS – G2056	Diagnosis of Idiopathic Environmental Intolerance

III. Indications and/or Limitations of Coverage

Application of coverage criteria is dependent upon an individual’s benefit coverage at the time of the request. Specifications pertaining to Medicare and Medicaid can be found in the “Applicable State and Federal Regulations” section of this policy document.

- 1) Specific IgE in-vitro allergy testing **MEETS COVERAGE CRITERIA** in **any** of the following situations:

- a) In lieu of skin testing for an **initial** allergy screen.
 - b) When skin testing is contraindicated (see Note 1).
 - c) When further treatment decisions would be impacted by confirmation of sensitivity in individuals for whom direct skin testing results are not consistent with the history of an anaphylactic or other severe reaction to an allergen.
- 2) When limited to allergens chosen for testing based on an individual's history, physical examination, and environment, specific IgE in-vitro allergy testing (up to 20 allergen specific antibodies per year) **MEETS COVERAGE CRITERIA**.
 - 3) In-vitro testing for total serum IgE **MEETS COVERAGE CRITERIA** in **any** of the following situations:
 - a) For individuals with moderate to severe asthma.
 - b) For individuals with signs or symptoms of allergic bronchopulmonary aspergillosis.
 - 4) To monitor for allergy resolution in children and adolescents with an initial positive food allergen result(s), annual re-testing for the same food allergen(s) **MEETS COVERAGE CRITERIA**.
 - 5) In the absence of a new clinical presentation, routine re-testing for allergies to the same allergens (except where specified above) **DOES NOT MEET COVERAGE CRITERIA**.
 - 6) The antigen leukocyte antibody test (ALCAT) **DOES NOT MEET COVERAGE CRITERIA**.

The following does not meet coverage criteria due to a lack of available published scientific literature confirming that the test(s) is/are required and beneficial for the diagnosis and treatment of an individual's illness.

- 7) For individuals with signs or symptoms of allergies, basophil activation flow cytometry testing (BAT) and in-vitro testing of IgG, IgA, IgM, **and/or** IgD **DO NOT MEET COVERAGE CRITERIA**.
- 8) In-vitro allergen testing using bead-based epitope assays (e.g., VeriMAP Peanut Dx) **DOES NOT MEET COVERAGE CRITERIA**.
- 9) For all situations, in-vitro testing using qualitative specific IgE multi-allergen screen that does not identify a specific allergen **DOES NOT MEET COVERAGE CRITERIA**.

NOTES:

Note 1: Skin testing is contraindicated in the following situations:

- Patients who have certain skin conditions (e.g., dermatographism, urticaria, cutaneous mastocytosis, atopic dermatitis, severe diffuse psoriasis)

- Patients who are taking medications that may interfere with the treatment of anaphylaxis (e.g., Beta-blockers and Angiotensin Converting Enzyme inhibitors) or may impair skin test sensitivity (e.g., tricyclic antidepressants, antihistamines)
- Patients who are at high risk to testing (e.g., poorly controlled asthma, clinical history of severe reaction to minute amounts of allergen, cardiac arrhythmia, unstable angina)
- Patients who have experienced an anaphylactic event within the past one month
- Uncooperative patients (e.g., small children, individuals with mental or physical impairments)

IV. Table of Terminology

Term	Definition
AAAAI	The American Academy of Allergy, Asthma, and Immunology
AAAI	Academy of Allergy, Asthma, and Immunology
AAFP	American Academy of Family Physicians
AAP	American Academy of Pediatrics
ACAAI	American College of Allergy, Asthma, and Immunology
ACD	Allergic contact dermatitis
ACR	American cockroach
AIT	Allergen immunotherapy
ALCAT	The Antigen Leukocyte Antibody Test
AR	Allergic rhinitis
Arah2-sIgE	<i>Arachis hypogaea</i> 2-specific immunoglobulin E
ARIA	Allergic Rhinitis and its Impact on Asthma
ATP	Atopy patch test
AUC	Area under the curve
BAT	Basophil activation flow cytometry testing
BAT	Basophil activation test
BBEA	Peanut bead-based epitope assay
CAA	Current allergic asthma
CAR	Current allergic rhinitis
CBS	Consensus based statements
CD4+	Cluster of differentiation 4
CLIA '88	Clinical Laboratory Improvement Amendments of 1988
CLSI	Clinical and Laboratory Standards Institute
CMS	Centers for Medicare & Medicaid Services
CRS	Chronic rhinosinusitis
CW	Choosing wisely
EAACI	European Academy of Allergy and Clinical Immunology
EDTA	Ethylenediaminetetraacetic Acid
EP	Expert panel
FA	Food allergy
FcεRI	High-affinity IgE receptor

FDA	The Food and Drug Administration
FEV1	Forced expiratory volume in 1 second
GA2LEN	Global Allergy and Asthma European Network
H1	H-1 receptor antagonists
H2	H-2 receptor antagonists
IgA	Immunoglobulin A
IgD	Immunoglobulin D
IgE	Immunoglobulin E
IgE-FAB	Immunoglobulin E- fragment antigen-binding region
IgG	Immunoglobulin G
IgG4	Immunoglobulin A
IgM	Immunoglobulin A
JTFPP	Joint Task Force on Practice Parameters
LCDs	Local coverage determinations
LDTs	Laboratory-developed tests
LEAP	Learning early about peanut allergy
MBB	Mucosal brush biopsies
MFI	Median fluorescence intensity
NASEM	The National Academies of Science, Engineering and Medicine
NIAID	National Institute of Allergy and Infectious Diseases
NICE	National Institute for Health and Care Excellence
NPV	Negative predictive value
NSAIDs	Nonsteroidal anti-inflammatory drugs
OFC	Oral food challenge
PAMD@	Precision allergy molecular diagnostic applications
PPV	Positive predictive value
RARS	Recurrent acute rhinosinusitis
SAR	Seasonal allergic rhinitis
sIgE	Specific immunoglobulin E
s-IgE	Specific immunoglobulin E
SPT	Skin prick tests
SSRIs	Selective serotonin reuptake inhibitors
ST	Skin test
Th2	T helper type 2
tIgE	Total immunoglobulin E
WAO	World Allergy Organization

V. Scientific Background

Allergies affect over 50 million Americans, including approximately 30 percent of adults and 40 percent of children (Jackson et al., 2013; NASEM, 2016). The incidence of allergic disease is increasing (Pawankar et al., 2013) and is estimated to result in over \$17 billion in health care costs and 200,000 emergency department visits annually (Adams et al., 2013).

A majority of environmental, food, and medication allergies with clinical significance are type I immunoglobulin E (IgE)-mediated allergies (Kowal & DuBuske, 2022). Diagnosis of an IgE-mediated allergy involves identification of the allergen, demonstration of IgE specific to that allergen, and confirmation that symptoms occur when the patient is exposed to the allergen. The IgE response to an allergen can be assessed using skin or serum testing. Patch testing is preferred for delayed T-cell mediated response (Chang & Guarderas, 2018; Zug et al., 2014).

Allergic diseases, respiratory infections, and autoimmune conditions have similar clinical presentations and self-reported symptoms have a relatively low positive predictive value (PPV) (Sampson et al., 2014). Thus, laboratory allergy and immunologic testing are useful in clarifying diagnosis and guiding treatment when the frequency, duration, and sequelae of upper respiratory infections exceed the norm or when rhinosinusitis or asthma symptoms persist despite treatment (Chow et al., 2012). Allergy testing is also useful in identifying causative allergen in atopic dermatitis (eczema), contact dermatitis, urticaria, angioedema, and food or drug allergies. Knowing the causal allergen helps provide clinically relevant information for avoidance and treatment (Chang & Guarderas, 2018).

Skin Testing

Skin testing is the most rapid, sensitive, and cost-effective testing modality for the detection of immunoglobulin E (IgE)-mediated disease. The procedure lasts less than an hour with minimal patient discomfort. There are several published practice parameters for allergen skin testing (Bernstein et al., 2008; Chang & Guarderas, 2018; Kowal & DuBuske, 2022).

Serum IgE

Immunoglobulin E (IgE) is one of five immunoglobulins and the one primarily involved in allergic disease. At the cellular level, the allergic response starts with “atopy,” a genetic predisposition to produce specific IgE after exposure to allergens. CD4+ helper T cells are predisposed to the “T helper type 2” (Th2) response, which causes the Th2 cells to secrete large amounts of interleukins 4 and 13, which then promotes production of the allergen specific IgE. From there, the allergen-specific IgE binds to high-affinity receptors on mast cells and basophils. At this point, if the relevant allergen is ingested in large enough amounts, the IgE molecules may cluster (cross-linking). This cross-linking causes the mast cells and basophils to release chemical and protein mediators, resulting in the characteristic allergic response (Stokes & Casale, 2022).

Immunoassays measuring both total IgE and allergen specific IgE in serum and other bodily fluids have been developed. Specific IgE immunoassays do not require patient cooperation, are not limited in patients with skin disease, are not blocked by antihistamines, and pose no risk of adverse reactions (Bernstein et al., 2008; Chang & Guarderas, 2018; Kowal & DuBuske, 2021; Stokes & Casale, 2022). Total IgE is usually unrelated to IgE levels for a specific allergen but may be useful in other conditions, such as asthma (Stokes & Casale, 2022).

Other testing

Patch testing is the gold standard for identification of a contact allergen (Mowad, 2006; Rietschel, 1997). Although occlusive patch testing is the most common technique, open, prophetic

(provocative), repeated insult, photopatch, and atopy patch tests are also available if special situations indicate their use (Bernstein et al., 2008).

Cellular activation assays measuring the release of histamine from basophils (Kim et al., 2016; Santos & Lack, 2016) or mast cells (Bahri et al., 2018) as diagnostic or prognostic indicators of allergy have been the subject of intense research. Basophil and eosinophilic reactivity tests have been found to be associated with food-induced allergic responses and have been shown in current research to be modified over time during immunotherapy (Sampson et al., 2014). In particular, the basophil activation test (BAT) has emerged as having superior specificity and comparable sensitivity to diagnose food allergies when compared with skin prick test and specific IgE (Santos & Shreffler, 2017). Histamine release from leukocytes of allergic persons is an excellent in vitro correlate of allergy; however, it is currently still considered a research test by the Academy of Allergy, Asthma, and Immunology (AAAAI) (Bernstein et al., 2008).

Basophil activation flow cytometry testing (BAT) has the potential to be a useful tool for measuring hypersensitivity to allergens, especially for patients who are not suitable for skin testing due to skin status or prior severe reactions since it is an ex vivo, flow cytometry-based assay. BAT, for use as standard clinical practice, is currently limited by its lack of standardization in methodology as well as between systems used. A study by Depince-Berger et al. (2017) has proposed standardization between systems and instruments using whole blood-ethylenediaminetetraacetic acid (EDTA) samples with instrumentation standardization. “BAT would strongly benefit from easy implementation [EDTA, one step stimulation/labeling, wash, full sample analysis over time parameter, B cell relative basophil count] and standardization of instrument settings on MFI [median fluorescence intensity] targets whatever system or instrument is used” (Depince-Berger et al., 2017). Hemmings et al. (2018) note that standardization, quality assurance, and clinical validation will facilitate the transition of the BAT from research to clinical practice.

Proprietary testing

The Antigen Leukocyte Antibody Test (ALCAT) is another test available for the assessment of allergens. ALCAT measures food/immune reactions through stimulation of leukocytes. The immunological reactions to this stimulation are intended to identify sensitivities regardless of pathway as antibodies do not necessarily need to be involved. Cell Science Systems suggests individuals with a variety of disorders (such as gastrointestinal, neurological, et al.) to take this test (Cell Science Systems, 2023). Although the ALCAT machine is FDA registered and there are a few papers published, results are not reproducible when subject to rigorous testing and do not correlate with clinical evidence of allergy (Beyer & Teuber, 2005; Hammond & Lieberman, 2018; Wuthrich, 2005).

Panels encompassing a large number of analytes are also offered by labs. For example, Genova Diagnostics offers a blood test for IgG and IgE antibodies for 87 different foods. Genova also offers several variations on this test, such as “Vegetarian” (21 foods), “Spices” (23 spices), “Molds” (15 molds), and more (Genova, 2023).

Spiriplex offers a microarray-style panel for allergen testing, called “Allergenex.” This test contains many purified allergen proteins to which a patient’s blood sample can bind. This binding

creates a quantifiable signal that allows the user to identify the number of IgE antibodies present, and therefore provide a picture of allergy. Spiriplex offers a test for 28 common food allergens, a test for 40 inhalant allergens, and 68 combined food and inhalant allergens (Spiriplex, 2023).

The VeriMAP Peanut Dx and the VeriMAP™ Peanut Sensitivity are both peanut-allergen specific bead-based epitope assays manufactured by AllerGenis LLC. According to AllerGenis, VeriMAP™ has a “95% positive predictive value and can reduce overdiagnosis and anxiety by minimizing false positives” (BioSpace, 2021). This is an emerging technology and additional peer-reviewed literature establishing the analytical validity, clinical validity, and clinical utility of such testing will be further required.

Analytical Validity

Variables that can influence the wheal size when performing skin prick tests (SPT) include multiple operators, extract concentrations and quality, skin test devices, time of day, location on the skin, and the measuring of results (Nelson, 2001; Werther et al., 2012).

In 2006, Oppenheimer and Nelson evaluated variability and analytical validity of skin testing. A questionnaire was sent to all physician and fellow members of the American College of Allergy, Asthma and Immunology who were currently practicing in the United States. The objective of this questionnaire was to determine the diversity of skin testing practices among allergists. The results showed great variability among physicians. In particular, “The average number of skin prick tests performed ranged from 5.09 (grasses) to 10.9 (trees), whereas the average number of intradermal tests performed ranged from 2.03 (grasses) to 5.6 (perennial). The allergen extract concentrations used for intradermal testing varied widely. Expressed as a dilution of the concentrated extracts, 20.8% use 1:100 dilutions, 10.3% use 1:500 dilutions, and 59.4% use 1:1,000 dilutions. Significant variability also occurred regarding devices and the technique with which the devices were used. Most clinicians (92.1%) used the most concentrated extract available for skin prick testing. For reporting the results of skin testing, 53.8% used a 0 to 4+ scale, and only 28.3% measured orthogonal diameters. Of those using a 0 to 4+ scale, two thirds related the results to the size of the histamine control” (Oppenheimer & Nelson, 2006). The results from this survey emphasize potential areas of improvement for allergists regarding skin test use and data.

The Clinical and Laboratory Standards Institute (CLSI) has evaluated the analytical validity of serum IgE measurements and found that “Clinical/diagnostic sensitivity and specificity of IgE antibody assays cannot be accurately determined due to the absence of definitive gold standard methods for defining allergic disease. Total and allergen-specific IgE analyses achieve among the highest analytical performance of any antibody assay by following consensus procedures in CLSI-ILA20-A3” (Hamilton et al., 2015).

Knight et al. (2018) “examined the qualitative concordance between SPT and sIgE as measured on the HYTEC™288 platform for 10 commonly encountered inhalant allergens”; a total of 232 subjects were included. Overall concordance between SPT and sIgE was >70% for all allergens tested. Sensitivity ranged from 25% to 95% depending on the allergen, while specificity was significantly higher for all allergens (78-97%). Negative predictive value (NPV) was >85% for all allergens tested, while PPV was more variable, ranging from 22% to 88%. The authors noted

that “these results are similar to findings in other studies comparing SPT with sIgE” (Knight et al., 2018).

Carlsson et al. (2015) examined the inter- and intra- variability of IgE and IgE receptor expression in the blood of seasonal allergic rhinitis (SAR) subjects. Thirty-two patients with SAR were included; the high-affinity IgE receptor, also known as FcεRI, and the low affinity receptor, also known as CD23, were measured. The authors found that “FcεRI expression on basophils and CD23 expression on B cells showed low intrasubject variability both in and out of the pollen season,” although there was a small seasonal difference with lower total IgE levels and FcεRI expression during the pollen season (Carlsson et al., 2015).

Siroux et al. (2017) explored the effect of allergen nature, route of exposure, and dose of exposure on IgE and IgG responses. A total of 340 patients (170 with asthma, 170 without) were included, and IgE/IgG responses to 47 inhalant and food allergens were analyzed and compared between five French regions according to route of allergen exposure (inhaled or food). “Ubiquitous” allergens (grass, olive/ash pollen, house dust mites) did not show marked difference in specific IgE level between regions. For region-specific allergens (ragweed, birch, cypress), IgE sensitization was associated with regional pollen exposure. Airborne allergens cross-reacting with food allergens led to frequent IgG recognition. The authors concluded that “the variability in allergen-specific IgE and IgG frequencies depends on exposure, route of exposure, and overall immunogenicity of the allergen. Allergen contact by the oral route might preferentially induce IgG responses” (Siroux et al., 2017).

Sookrung et al. (2019) measured the agreement of a SPT and serum specific IgE test to *Periplaneta americana* (American cockroach, ACR) allergies. ACR-extract was used, and sera was obtained from 66 individuals clinically diagnosed with chronic allergic rhinitis. Of the 66 samples, 46 were positive and 20 negative after a SPT to ACR-extract. Serum IgE levels were then measured by a commercial test kit. The authors note that of the SPT positive cases to ACR-extract, only 32.6% were also positive for serum IgE, indicating low concordance between the two testing methods (Sookrung et al., 2019).

He and Reisacher (2019) measured the sensitivity, specificity, and predictive value of oral mucosal brush biopsies (MBB) as a new diagnostic test for peanut allergies. Twenty individuals participated in this study; each participant underwent oral MBB and serum testing for peanut IgE. The authors note that “At 0.12 kU/L, the sensitivity of oral MBB testing was 80% and the specificity was 85%, whereas at 1.0 kU/L, the sensitivity of sIgE testing was 50% and the specificity was 100%. From the ROC curves, the areas under the ROC curve (AUC) for oral MBB and sIgE were 0.91 ($p < 0.001$) and 0.74 ($p = 0.007$), respectively. Combination testing further increased both sensitivity and accuracy over oral MBB alone” (He & Reisacher, 2019). These results are promising for oral MBB, although more research needs to be completed.

Clinical Utility and Validity

In 1998, Tschopp et al. (1998) compared three diagnostic tests for atopic diseases. Total serum IgE, Phadiatop, and the SPT were compared for 8329 individuals. Current allergic asthma (CAA) and current allergic rhinitis (CAR) were the conditions studied. The prevalence of CAA was 1.8% and prevalence for CAR was 16.3%. The prevalence of positive tests was 29%, 23%,

and 23% for Phadiatop, SPT, and IgE, respectively. The results were as follows: “To diagnose current allergic asthma (CAA) and current allergic rhinitis (CAR), the sensitivity of Phadiatop was significantly higher than that of SPT (72.5% vs 65.4%, 77.1% vs 68.4% respectively) and IgE (72.5% vs 56.9%, 77.1% vs 43.9%, respectively). The sensitivity of SPT was significantly higher (68.4% vs 43.9%) than that of IgE to diagnose CAR. When CAA and CAR were excluded, the SPT specificity was significantly higher than that of Phadiatop (77.8% vs 71.9% and 85.9% vs 80.5%, respectively): when CAR was excluded, SPT was significantly higher than IgE (85.9% vs 81.4%). SPT had significantly the best positive predictive value for CAA (5.2% for SPT vs 4.6% for both IgE and Phadiatop) and CAR (48.7% for SPT vs 43.5% for Phadiatop and 31.6% for IgE). The three markers of atopy had roughly the same negative predictive value (NPV) for CAA, but IgE had a significantly lower NPV for CAR than SPT and Phadiatop (88.1% vs 93.3% and 94.7%, respectively). The diagnostic efficiency of SPT was significantly higher than that of Phadiatop (83.1% vs 79.9% and 77.6 vs 71.9%, respectively) to diagnose CAR and CAA. IgE and SPT had equal efficiency (77.6%), which was significantly higher than that of Phadiatop, to diagnose CAA (71.9%)” (Tschopp et al., 1998). The authors concluded that “SPT have the best positive predictive value and the best efficiency to diagnose respiratory atopic diseases. Furthermore, SPT give information on sensitivity to individual allergens and should therefore be used primarily by clinicians to assess respiratory allergic diseases” (Tschopp et al., 1998).

A retrospective analysis included patients who had been prick tested to “establish whether an incomplete diagnosis would have been reached if patch testing had been omitted.” The authors observed that if “investigation of allergic skin disease is undertaken by a non-dermatologist, it is unlikely that patch testing will be performed.” A total of 330 patients had been prick tested in the time period specified. Sixty-eight patients had positive reactions on prick testing, and 36 of those had positive patch tests. Of the 262 patients who had negative prick tests, 121 had positive patch tests (46.1%) of current relevance to patient history in 92 subjects (35.1%). The authors concluded that “omission of patch testing from the investigation of allergic skin disease, even when contact urticaria may be the sole suspected diagnosis, would result in the frequent missed diagnosis of contact allergy” (Usmani & Wilkinson, 2007).

In 2014, a meta-analysis examined the clinical validity of SPT and IgE measurement for food allergy. Twenty-four studies consisting of 2831 participants were included. The results were as follows: “For cows' milk allergy, the pooled sensitivities were 88% (SPT), and 87% (IgE) and specificities were 68% and 48%. For egg, pooled sensitivities were 92% and 93% and specificities were 58% and 49% for SPT and specific-IgE. For wheat, pooled sensitivities were 73% and 83% and specificities were 73% and 43% for SPT and sIgE. For soy, pooled sensitivities were 55% and 83% and specificities were 68% and 38% for SPT and sIgE. For peanut, pooled sensitivities were 95% and 96%, and specificities were 61% and 59% for SPT and sIgE” (Soares-Weiser et al., 2014).

Klemans et al. (2015) examined the diagnostic accuracy of using sIgE to peanut components to improve sensitivity and specificity of peanut allergen testing. Twenty-two studies were included. The authors found that “sIgE to Ara h 2 [a peanut component] showed the best diagnostic accuracy of all diagnostic tests to diagnose peanut allergy. Compared to the currently used SPT and sIgE to peanut extract, sIgE to Ara h 2 was superior in diagnosing peanut allergy” (Klemans et al., 2015). The authors also found that the worst accuracy was observed to be sIgE to Ara8 and

Ara9. The authors concluded that “sIgE to Ara 2 should replace SPT and sIgE to peanut extract in daily clinical practice” (Klemans et al., 2015).

Caglayan Sozmen et al. (2015) examined the diagnostic accuracy of using the patch test to avoid oral food challenge (OFC). They found that in 243 children that underwent OFC to suspected food, clinically relevant food allergies were seen in 40 (65%) children to egg and in 22 (35%) to cow's milk. The sensitivity of SPT for both milk and egg were 92%, specificity 91%, PPV 35%, and NPV 93%. Sensitivity, specificity, PPV, and NPV of atopy patch test for both milk and egg were 21%, 73%, 20%, and 74%, respectively.

Santos et al. (2014) studied the performance of basophil activation tests (BAT) as a diagnostic marker for peanut allergy. Forty-three peanut-allergic children, 36 peanut-sensitized but tolerant children, and 25 non-peanut-sensitized nonallergic children underwent SPT, sIgE, and BAT. The authors found that BAT in peanut-allergic children showed a peanut dose-dependent upregulation of CD63 and CD203c while there was no significant response in the other two cohorts. BAT optimal diagnostic cutoffs showed 97% accuracy, 95% PPV, and 98% NPV. BAT allowed reduction of required oral food challenges (OFCs) by two-thirds. BAT proved particularly useful in cases in which specialists could not accurately diagnose peanut allergy with SPT and sIgE to peanut and to AraH2. Using a two-step diagnostic approach in which BAT was performed only after equivocal SPT or AraH2-sIgE, BAT had a major effect (97% reduction) on the number of OFCs required.

Santos et al. (2015) also studied the utility of BAT to predict the severity and reactivity to peanut during OFCs. They found that “Of the 124 children submitted to OFCs to peanut, 52 reacted with clinical symptoms that ranged from mild oral symptoms to anaphylaxis. Severe reactions occurred in 41% of cases, and 57% reacted to 0.1 g or less of peanut protein. The ratio of the percentage of CD63(+) basophils after stimulation with peanut and after stimulation with anti-IgE (CD63 peanut/anti-IgE) was independently associated with severity, whereas the basophil allergen threshold sensitivity CD-sens ($1/EC_{50} \times 100$, where EC_{50} = half maximal effective concentration) value was independently associated with the threshold of allergic reactions to peanut during OFCs. Patients with CD63 peanut/anti-IgE levels of 1.3 or greater had an increased risk of severe reactions (relative risk, 3.4). Patients with a CD-sens value of 84 or greater had an increased risk of reacting to 0.1 g or less of peanut protein (relative risk, 1.9)” (Santos et al., 2015). The authors concluded that “Basophil reactivity is associated with severity, and basophil sensitivity is associated with the threshold of allergic reactions to peanut. CD63 peanut/anti-IgE and CD-sens values can be used to estimate the severity and threshold of allergic reactions during OFCs” (Santos et al., 2015).

Davila et al. (2015) explored the association between total IgE and severity of asthma. A total of 383 patients were included (129 mild, 82 moderate, and 172 severe). Serum IgE levels were noted to vary “markedly” (147% coefficient of variation). The authors did not find an association between total IgE and forced expiratory volume in one second (FEV1) or asthma severity; although, the severe subgroup had a higher percentage of patients with >400 IU/mL. Independent predictors of higher IgE were found to be younger age, sensitization to at least two allergens, male gender, and family history of asthma. The authors concluded that “we did not find a significant association between serum total IgE levels and asthma severity or airflow limitation,

except for a higher percentage of patients with IgE > 400 IU/mL in the severe subgroup” (Davila et al., 2015).

Tannert et al. (2017) investigated the relevance of a positive skin test and positive IgE test to penicillin allergy. Twenty-five patients with positive results were given penicillin, and another 19 patients deemed allergic were included. However, only nine of the 25 patients given penicillin were challenge positive. Positive results from each test alone did not predict allergy. The authors concluded that “the best predictor for a clinically significant (IgE-mediated) penicillin allergy is a combination of a positive case history with simultaneous positive ST result and s-IgE or a positive challenge result” (Tannert et al., 2017).

Suárez-Fariñas et al. (2021) investigated the validity of the peanut BBEA diagnostic test on 133 subjects as well as on 82 additional subjects from another study, forming a cohort for a paper titled, “Accurate and reproducible diagnosis of peanut allergy using epitope mapping.” The authors measured levels of IgE to epitopes evaluated against a threshold established prior to the study. The peanut BBEA diagnostic test diagnosed 93% of subjects accurately, with a sensitivity threshold of 92% and specificity of 94%. The positive predictive value (PPV) was 91%. The authors concluded that “the overall accuracy was found to be superior to existing diagnostic tests for peanut allergy including skin prick testing, peanut sIgE, and peanut component sIgE testing” (Suárez-Fariñas et al., 2021).

VI. Guidelines and Recommendations

The American Academy of Allergy, Asthma, and Immunology (AAAAI) and the American College of Allergy, Asthma, and Immunology (ACAAI)

The AAAAI and ACAAI published practice parameters in 2008 for allergy testing (Bernstein et al., 2008) which noted that “For individual patients, the choice of test allergens is guided by the history and physical examination and the physician’s knowledge, training, and experience.” The guidelines recommended that “Specific IgE immunoassays may be preferable to skin testing under special clinical conditions, such as widespread skin disease, patients receiving skin test suppressive therapy, uncooperative patients, or when the history suggests an unusually greater risk of anaphylaxis from skin testing.” They also note that for both skin testing and in-vitro specific IgE testing, “the allergens selected ... should be determined based on the patient’s age, history, environment and living conditions (e.g., region of the country), occupation, and activities.” Also, “The best indicators in the selection of appropriate pollens for clinical use are extensive prevalence in the air and concurrent allergy symptoms during annually recurrent seasons when such pollens are expected to be present in the ambient air.”

The AAAAI and ACAAI guidelines also state, “As is the case with skin tests, a direct correlation cannot be assumed between the presence of specific IgE (sIgE) antibodies and clinical disease.” Additionally, “sensitivity and the positive predictive value of both prick/puncture and specific IgE tests generally tend to be higher among pollens, stable anaphylactogenic foods, house dust mite, certain epidermals, and fungi compared with venoms, drugs, and chemicals.”

With regards to total IgE testing, these groups indicate, “Measurements of total serum IgE concentration are of modest clinical value when used as a screen for allergic disease or for predicting the risk of allergic disease.”

The AAAAI and ACAAI also note that “IgG and IgG subclass antibody tests for food allergy do not have clinical relevance, are not validated, lack sufficient quality control, and should not be performed.”

Regarding basophil activation assays they state, “Histamine and leukotriene release measurements from human basophils after incubation with allergen are valuable research tools for in vitro investigations of allergy” (Bernstein et al., 2008).

Their practice parameter on drug allergy also states that “The basophil activation test is a recently described method of evaluating expression of CD63 on basophils after stimulation with an allergen. There are limited data using this method to evaluate patients with possible allergies to β -lactam antibiotics and nonsteroidal anti-inflammatory drugs (NSAIDs)” (Boyce et al., 2010).

They also recommend, “Because anaphylactic reactions cannot be distinguished from anaphylactoid, nonimmune occurrences, it has been recommended that plasma histamine, tryptase, and specific IgEs (if available) may be ordered at the time of reaction and skin tests be performed later” (Boyce et al., 2010).

In their 2014 practice parameter on food allergy (Sampson et al., 2014) they acknowledge: “Basophil and eosinophilic reactivity tests have been shown to be associated with food-induced allergic responses and have been shown in current research to be modified over time during immunotherapy.”

Their 2014 practice parameter on rhinosinusitis also recommends to “Perform an evaluation for specific IgE antibodies to airborne allergens in patients with RARS or CRS.” An updated practice parameter on rhinitis published in 2020 comments that local allergic rhinitis will often be associated with “negative skin prick tests (and intradermal tests, when performed) and absence of serum-specific IgE (sIgE) antibodies but a positive nasal allergen provocation test (NAPT) to aeroallergens (Dykewicz et al., 2020). With respect to vasomotor rhinitis, the authors state that “laboratory tests, skin prick tests, and sIgE are helpful only to exclude AR [allergic rhinitis]” (Dykewicz et al., 2020).

In this practice parameter, they also make the following summary concerning re-evaluation of food allergies in children and adolescents: “Summary Statement 11: Consider the natural course of allergies to specific foods when deciding on the frequency of food allergy follow-up evaluations, recognizing that allergies to certain foods (milk, egg, wheat, and soy) generally resolve more quickly in childhood than others (peanut, tree nuts, fish, and shellfish). These observations could support individualized follow-up (ie, roughly yearly re-evaluations of these allergies in childhood) with less frequent retesting if results remain particularly high (eg, >20-50 kUA/L). [Strength of recommendation: Moderate; C Evidence]” (Dykewicz et al., 2020).

In their 2015 practice parameter on anaphylaxis (Lieberman et al., 2015), they recommend “Skin tests and/or in vitro tests for specific IgE and challenge tests might be appropriate to help define the cause of the anaphylaxis.”

They also recommend against routinely obtaining total serum IgE levels for the diagnosis of food allergy, however, because of the low PPV of self-reported symptoms and lack of pathognomonic signs on physical examination, they recommend that the accurate diagnosis of IgE-mediated food allergy should be aided by laboratory allergy testing, including skin prick and/or serum IgE testing. The clinician should use specific IgE tests (skin prick tests, serum tests, or both) to foods as diagnostic tools; however, testing should be focused on foods suspected of provoking the reaction, and test results alone should not be considered diagnostic of food allergy. Moreover, “the diagnosis of food-induced anaphylaxis should be based on signs and symptoms in association with likely or known exposure to a food allergen”, as “events mimicking anaphylaxis also can occur after the ingestion of food” (Lieberman et al., 2015).

In a Choosing Wisely (CW) report, the AAAAI recommends against performing “unproven diagnostic tests, such as immunoglobulin G (IgG) testing or an indiscriminate battery of immunoglobulin E (IgE) tests, in the evaluation of allergy” (AAAAI, 2012).

In another CW report, the AAAAI recommends against routine diagnostic testing in patients with chronic urticaria, stating that “skin or serum-specific IgE testing for inhalants or foods is not indicated, unless there is a clear history implicating an allergen as a provoking or perpetuating factor for urticaria” (AAAAI, 2012).

The AAAAI also published a 2020 practice parameter update on peanut allergy diagnosis. The authors recommend in favor of diagnostic skin prick test or sIgE testing for peanut allergy in patients with physician-judged high pretest probability of peanut allergy. Testing is also recommended prior to an oral food challenge for patients with moderate pretest probability of peanut allergy. Ara h 2 diagnostic testing is the suggested approach for patients presenting for evaluation of suspected peanut allergy for which a single diagnostic test is to be used, due to its superior diagnostic accuracy “by virtue of more optimal positive/negative likelihood ratios.” However, Ara h 2 is noted to have lower sensitivity than the skin prick or sIgE tests, so a clinician may use Ara h 2, SPT, or sIgE to confirm the diagnosis of peanut allergy in a patient with a high prior probability. The AAAAI recommends against “routine use of component testing in addition to either SPT or sIgE to whole peanut to increase diagnostic accuracy,” and against using the results of skin prick or sIgE to determine “the severity of an allergy phenotype or to predict the severity of a future reaction” (Greenhawt et al., 2020). It is noteworthy that all the recommendations above were assigned “low” or “very low” degrees of evidence certainty (Greenhawt et al., 2020).

Joint Task Force on Practice Parameters (JTFPP)

In a practice parameter concerning contact dermatitis, the Joint Task Force on Practice Parameters—composed of the American Academy of Allergy, Asthma & Immunology (AAAAI), the American College of Allergy, Asthma & Immunology (ACAAI), and the Joint Council of Allergy, Asthma & Immunology—proposed this series of summary statements:

“Summary Statement 1: Consider ACD [allergic contact dermatitis] in the differential diagnosis of patients with chronic eczematous or noneczematous dermatitis. [Strength of Recommendation: Strong; C Evidence]

Summary Statement 2: In patients suspected of ACD, patch testing is the gold standard to confirm the diagnosis. [Strength of Recommendation: Strong; C Evidence]

Summary Statement 3: In addition to personal products used by a patient suspected of ACD, review the home and workplace for other sources of contact allergens. [Strength of Recommendation: Moderate; D Evidence] Summary Statement

Summary Statement 4: Evaluate patients for both irritant and allergic causes, especially in those presenting with hand dermatitis. [Strength of Recommendation: Strong; C Evidence]

Summary Statement 5: Allergic CD should be suspected and evaluated in the patient with both generalized and anatomically localized skin eruptions (such as the hands, face, eyelids) that come in contact with the substances in the environment. [Strength of Recommendation: Moderate; C Evidence]

Summary Statement 6: In a patient with a facial rash involving the periorbital areas (eg, eyelids), evaluate for ACD caused by components of cosmetics, such as fragrances, preservatives, and excipients, because these are common sensitizers of the facial skin. [Strength of Recommendation: Moderate; C Evidence]

Summary Statement 7: Evaluate patients presenting with lip dermatitis (cheilitis) and perioral dermatitis for both irritant and allergic causes of contact dermatitis. [Strength of Recommendation: Moderate; C Evidence]

Summary Statement 8: Evaluate patients with chronic oral mucosal inflammatory conditions for disorders other than ACD. [Strength of Recommendation: Moderate; C Evidence]

Summary Statement 9: In patients presenting with dermatitis that involves the scalp and neck, consider patch testing for common causative sensitizers in cosmetics, hair products, and jewelry. [Strength of Recommendation: Moderate; C Evidence]

Summary Statement 10: Consider irritant and ACD in all patients presenting with acute or chronic hand eczema. All such patients suspected of CD should undergo patch testing. [Strength of Recommendation: Moderate; C Evidence]

Summary Statement 11: Evaluate patients with axillary dermatitis for ACD caused by local contact sensitivity to allergens in topically applied products found in deodorants and textiles. In some cases, axillary dermatitis could be a manifestation of systemic contact dermatitis (SCD) (i.e., “the baboon syndrome”). [Strength of Recommendation: Moderate; C Evidence]

Summary Statement 12: Evaluate patients presenting with anogenital dermatitis for possible ACD to antigens contained in topically applied products. [Strength of Recommendation: Moderate; C Evidence]

Summary Statement 13: Consider a diagnosis of SCD following systemic exposure (e.g., ingestion, infusion, or transcutaneous exposure) to a known contact sensitizer in a patient who presents with generalized dermatitis, intertriginous and flexural exanthema (Baboon syndrome), and/or a flare at previous cutaneous sites of exposure [Strength of Recommendation: Moderate; C Evidence].

Summary Statement 14: Consider PT to rubber chemicals, adhesives, and leather components of footwear in patients presenting with unexplained chronic dermatitis involving the lower extremities, feet and/or soles. [Strength of Recommendation: Moderate; C Evidence]

Summary Statement 15: In addition to avoiding irritants in patients with atopic dermatitis (AD), evaluate for ACD, if suspected, as the 2 dermatologic conditions often coexist in the same patient. [Strength of Recommendation: Moderate; C Evidence]” (Fonacier et al., 2015).

Consensus based statements (CBSs) regarding the diagnosis and management of rhinitis from the JTFPP include the following (Dykewicz et al., 2020):

Recommendation no.	CBS or GRADE recommendation	Strength of recommendation	Certainty of evidence
1	CBS: We recommend that the clinician complete a detailed history and a physical examination in a patient presenting with symptoms of rhinitis.	Strong	Low
2	CBS: We recommend that for patients presenting with rhinitis symptoms, a review of all current medications should be completed to assess whether drug-induced rhinitis may be present.	Strong	Ungraded
3	CBS: We recommend that aeroallergen skin prick testing or sIgE testing be completed to confirm the diagnosis of AR in a patient with a history consistent with AR.	Strong	High
4	CBS: We recommend that the clinician not perform food skin prick testing or sIgE for foods in their routine evaluation of a patient presenting with the signs and symptoms compatible with the diagnosis of AR.	Strong	Ungraded
5	CBS: We suggest that the use of a validated instrument (eg, scoring system, scale, or questionnaire) be considered to help determine the severity of rhinitis and to monitor the degree of disease control.	Conditional	Low

World Allergy Organization Position Paper

In 2020, the World Allergy Organization published a position paper on IgE allergy diagnostics and other relevant allergy tests. Key statements from the paper can be found below:

- “Clinical suspicion of allergic sensitization is confirmed by demonstrating the presence of allergen-specific IgE antibodies *in vivo* (skin tests) or *in vitro*.
- Confirmation of allergen sensitization and the identification of causal allergens are essential for optimizing the management of allergic conditions.
- Skin prick testing (SPT) is the most frequently used method for the detection of IgE antibodies, due to its rapidity, simplicity, and low cost. Skin prick tests and other skin test results must be interpreted by a clinician with adequate knowledge of medical history, clinical findings, and relevant type I allergens (including environmental, food, animal, insect, fungal, and drug allergens). Skin tests should include the relevant allergens in the given geographical area and ideally carried out only using standardized allergenic extracts.
- *In vitro* tests, including molecular based allergy diagnostics, using either in single-plex and in multi-plexed strategies and other more functional tests, such as Basophil Activation Tests allow to better define the IgE profile of the patient. This approach is in line with the Precision Medicine statements” (Ansotegui, Melioli, Canonica, Caraballo, et al., 2020)

The paper also states that “Skin tests, especially SPT, represent the most reliable and cost-effective tool for the diagnosis and management of IgE-mediated diseases. They demonstrate a good correlation with outcomes of nasal, conjunctival, dermal, oral, and bronchial challenges” (Ansotegui, Melioli, Canonica, Caraballo, et al., 2020).

Clinical conditions where SPT is indicated include:

- “Asthma;
- Rhinitis/rhinosinusitis/rhino-conjunctivitis/conjunctivitis;
- Eczema/atopic dermatitis (in the setting of selectively high clinical suspicion for underlying presence of IgE hypersensitivity to specific allergens);
- Suspected food allergy (oral allergy syndrome, anaphylaxis/acute onset or exacerbation of urticaria or eczema that is temporally correlated with food ingestion);
- Suspected drug allergy;
- Hymenoptera venom allergy (systemic reactions immediately following insect sting);
- Suspected occupational disease or exposure to selected potential allergens;
- Chronic urticaria in rare selected cases which strongly suggest an allergen as potential trigger/ aggravating factor;
- Less common disorders, such as eosinophilic esophagitis, eosinophilic gastroenteritis, or allergic bronchopulmonary aspergillosis, where IgE sensitization is one of the characteristics of its pathogenesis. However, there is controversy regarding the utility of SPT for these illnesses” (Ansotegui, Melioli, Canonica, Caraballo, et al., 2020)

“SPT is not routinely indicated in the following instances in the absence of other existing features of allergic disease:

- Suspected food intolerance (e.g., irritable bowel syndrome, etc.);
- Chronic urticaria in the absence of allergic features in the history;
- Desire to lose weight (according to nonconventional approaches, obesity may be due to food intolerance, but no supporting scientific data have been reported in the literature);
- Non-specific food-associated symptoms to food additives/preservatives/colorants;
- Evaluation of the effectiveness of allergen immunotherapy (but may be supportive in Hymenoptera venom immunotherapy);
- Non-specific respiratory symptoms to irritants (i.e., smoke, perfumes, detergents, chemicals and other strong odors);
- Screening for allergic sensitization patterns in the absence of clinical symptoms (i.e., family history of allergy);
- Non-specific cutaneous rashes in the absence of atopic features or other allergic symptoms; migraine, except for the indication of specific hypersensitivity to hormones. However, strong scientific data are still missing.
- Chronic fatigue syndrome” (Ansotegui, Melioli, Canonica, Caraballo, et al., 2020)

In a 2020 publication on anaphylaxis guidance, the WAO confirms that “allergy testing should be based on patient history and local data regarding the common causes of anaphylaxis in the region. The most frequent elicitor groups worldwide are food, insect venom, and drugs” (Cardona et al., 2020).

World Allergy Organization (WAO), Allergic Rhinitis and its Impact on Asthma (ARIA), and the Global Allergy and Asthma European Network (GA²LEN)

The WAO, ARIA, and GA²LEN published a consensus document in 2020 focused on molecular-based allergy diagnoses. Precision allergy molecular diagnostic applications (PAMD@) “can increase the accuracy of an allergy diagnosis in certain circumstances. In allergic patients, a molecular approach is suitable for the following:

- Assessing the risk of potential allergic reactions, which depend on the individual allergic (clinical) sensitization profile;
- Evaluating whether unknown potential triggering factors are present (i.e., the presence of sIgE versus allergenic molecules correlated with high risk for allergic reactions)” (Ansotegui, Melioli, Canonica, Gomez, et al., 2020).

National Institute of Allergy and Infectious Diseases (NIAID)

The NIAID convened an expert panel to review current information and to make recommendations related to the evaluation of food allergy (FA), including the use of specific IgE (sIgE) testing (Boyce et al., 2010). With regards to allergen-specific serum IgE determination, NIAID recommended that “sIgE tests for identifying foods that potentially provoke IgE-mediated food-induced allergic reactions, but alone these tests are not diagnostic of FA.” It stated that “sIgE testing and skin prick testing both depend on the presence of allergen-specific antibodies. Because the former test measures sIgE in the serum and the latter reflects IgE bound to cutaneous mast cells, their results may not always correlate. Serum testing can be especially useful when SPTs cannot be done (for example, due to extensive dermatitis or dermatographism), or when antihistamines cannot be discontinued.” The NIAID also recommended not using the combination of skin prick test (SPT), sIgE tests and atopy patch test (ATP) for the routine diagnosis of food allergy.

Additionally, the NIAID notes that “the routine use of measuring total serum IgE should not be used to make a diagnosis of FA” (Boyce et al., 2010).

“Non-standardized tests” such as basophil histamine release/activation, lymphocyte stimulation, allergen-specific IgG, cytotoxicity assays, and mediator release assays should not be used in the routine evaluation of FA, according to the NIAID guidelines (Boyce et al., 2010).

In 2017, the NIAID published addendum guidelines for the prevention of peanut allergy in the United States. These guidelines note that the expert panel (EP) “recommends that evaluation with peanut-specific IgE (peanut sIgE) measurement, SPTs, or both be strongly considered before introduction of peanut to determine if peanut should be introduced and, if so, the preferred method of introduction. To minimize a delay in peanut introduction for children who may test negative, testing for peanut sIgE may be the preferred initial approach in certain health care settings, such as family medicine, pediatrics, or dermatology practices, in which skin prick testing is not routine” (Togias et al., 2017). Further, “The EP does not recommend food allergen panel testing or the addition of sIgE testing for foods other than peanut because of their poor positive predictive value, which could lead to misinterpretation, overdiagnosis of food allergy, and unnecessary dietary restrictions” (Togias et al., 2017). More, if an infant has severe eczema, an

egg allergy, or both, the EP recommends to “Strongly consider evaluation by sIgE measurement and/or SPT and, if necessary, an OFC. Based on test results, introduce peanut-containing foods” (Togias et al., 2017).

American Academy of Pediatrics (AAP)

In 2012, AAP released a clinical report on allergy testing in childhood. It stated that “Both serum sIgE tests and SPT are sensitive and have similar diagnostic properties.” The AAP summary included the following:

- “Treatment decisions for infants and children with allergy should be made on the basis of history and, when appropriate, identified through directed serum sIgE or SPT testing. Newer in vitro sIgE tests have supplanted radioallergosorbent tests.”
- “Positive sIgE test results indicate sensitization but are not equivalent to clinical allergy. Large panels of indiscriminately performed screening tests may, therefore, provide misleading information.”
- “Increasingly higher levels of sIgE (higher concentrations on serum tests or SPT wheal size) generally correlate with an increased risk of clinical allergy.”
- “Use of a multiallergen serum test can be helpful for screening for atopic disease if there is a clinical suspicion. If positive, allergen-specific testing may be considered.
- “Tests for allergen-specific IgG antibodies are not helpful for diagnosing allergies” (AAP, 2012).

In 2019, the AAP published new guidelines on the prevention of childhood food allergies and other allergic conditions. This article states that “The new recommendations for the prevention of peanut allergy are based largely on the LEAP trial and are endorsed by the AAP.” The AAP endorsed guidelines were published by Togias et al. (2017) and are noted above. They state that the highest-risk infants (those with severe eczema and/or egg allergies) should be introduced to peanuts by four to six months; further, allergy testing is strongly advised before peanut introduction. SPT and blood testing for peanut-specific IgE (sIgE) are allowable (Greer et al., 2019; Sicherer, 2017).

In 2020, the AAP published a state-of-the-art review of peanut allergy testing advances and controversies. The article states that “current first-line diagnostic tests for peanut allergy have limited specificity, which may be enhanced with emerging tools such as component-resolved diagnostics.” Like the 2019 guideline, they note that first-line best practices for peanut allergy testing include SPT or serum peanut-specific IgE measurement. While both tests are highly sensitive, neither correlate strongly with reaction severity, according to the AAP (Abrams et al., 2020).

U.S. Food and Drug Administration (FDA) on Xolair

The availability of Xolair for treatment of allergic asthma also has implications for allergy testing. According to the package insert, Xolair is indicated for patients six years of age and older with moderate to severe persistent asthma who have a positive skin test or in vitro reactivity to a perennial aeroallergen and whose symptoms are inadequately controlled with inhaled

corticosteroids... Determine dose (mg) and dosing frequency by serum total IgE level (IU/mL) measured before the start of treatment, and by body weight (kg).” The prescribing information also notes that “Total IgE levels are elevated during treatment and remain elevated for up to one year after the discontinuation of treatment. Therefore, re-testing of IgE levels during Xolair treatment cannot be used as a guide for dose determination” (FDA, 2016).

International Consensus Statement on Allergy and Rhinology: Allergic Rhinitis

The authors reviewed the existing evidence behind various aspects of evaluation and diagnosis of the AR patient, and developed the following recommendations for AR diagnostic modalities (Wise et al., 2023):

- Patient history: "Using history to make a presumptive diagnosis of AR is reasonable and would not delay treatment initiation. History should be combined with physical examination, which may not be possible in some scenarios such as telemedicine. Confirmation with diagnostic testing is required for progression to AIT or targeted avoidance therapy, or desirable with inadequate response to treatment... Despite low level evidence specifically addressing this area, history is essential in the diagnosis of AR."
- Physical examination: "When possible, physical examination should be performed with appropriate personal protective equipment to aid in the diagnosis of AR and exclusion of other conditions. When combined with patient history, it increases diagnostic accuracy and may exclude alternative causes of symptoms."
- Nasal endoscopy: "Nasal endoscopy may be considered as a diagnostic adjunct in the evaluation of patients with suspected AR."
- Radiologic studies: "Routine use of imaging is not recommended for the diagnosis of AR."
- Use of validated subjective instruments and patient-reported outcome measures: "Validated surveys may be used to screen for AR, follow treatment outcomes and as a primary outcome measure for clinical trials. Specific tests are optimized for various clinicopathological scenarios."
- Skin prick testing: "Patients can benefit from identification of their specific sensitivities. Skin prick testing (SPT) is a quick and relatively comfortable way to test several antigens with accuracy similar to other available methods of testing... Regular use of the same SPT device type will allow clinicians to familiarize themselves with it and interpretation of results may therefore be more consistent. The use of standardized allergen extracts can further improve consistency of interpretation."
- Skin intradermal testing: "Intradermal skin tests may not perform as well as SPT in most clinical situations... Intradermal testing may be used to determine aeroallergen sensitization in individuals suspected of having AR."
- Blended skin testing techniques: "While AIT can be based off SPT results alone, endpoint-based immunotherapy may have possible benefits of decreased time to therapeutic dosage... Blended skin testing techniques, such as modified quantitative testing, are methods that can be used to determine a starting point for AIT or confirm allergic sensitization."
- Serum total immunoglobulin E: "Assessment of total IgE may be useful to assess overall atopic status; furthermore, in selected cases it might help guide therapy (i.e., predict outcome of AIT)."

- Serum allergen-specific immunoglobulin E: “Patients can benefit from identification of their specific sensitivities. Further, in some patients who cannot undergo SPT, serum sIgE testing is a safe and effective alternative... Serum sIgE testing may be used in patients who cannot undergo allergy skin testing. The use of highly purified allergen or recombinants can increase the sensitivity, specificity, and diagnostic accuracy of sIgE tests. Rigorous proficiency testing on the part of laboratories may also improve accuracy.”
- Nasal allergen-specific immunoglobulin E: “In patients with non-allergic rhinitis who also have risk factors for atopic disease and have inadequate response to pharmacotherapy, testing for nasal sIgE may be helpful in confirming a diagnosis of local AR and allowing for treatment with AIT. There is no consensus for levels of nasal sIgE that indicate sensitivity... Measurement of nasal sIgE is an option in patients with non-allergic rhinitis suspected of having local AR to support this diagnosis and guide AIT if pharmacologic therapies are inadequate. Consensus for levels of nasal sIgE indicating AR need to be established.”
- Basophil activation test: “The evidence does not support routine use for the diagnosis of AR or for following AIT response... Application of basophil activation test in specific situations where other diagnostic procedures for AR are not possible or conflicting. Potentially useful for monitoring AIT if other methods fail or show conflicting results.”
- Component resolved diagnostic testing: “Molecular diagnosis may be a useful tool for assessment of AR in some scenarios, especially in polysensitized patients... Component resolved diagnostic testing is an option for diagnosis of AR by specialists.”
- Nasal provocation testing: “Application of nasal provocation testing is useful in local AR and to confirm occupational rhinitis.”
- Nasal cytology: “Nasal cytology could help in cases of non-allergic rhinitis to suspect local AR or in cases of AR to diagnose a mixed rhinitis. It could be considered an option in cases of negative SPT and/or serum sIgE to evaluate the presence of mucosal eosinophils and consideration of local AR or type two inflammation. The cut-off values for determining non-allergic rhinitis with eosinophilia syndrome (NARES) are not yet clear.”
- Nasal histology: “Nasal histology may be helpful in clinical research or selected cases (e.g., evaluation of tissue eosinophils during surgery). Recommendation against in routine clinical practice for AR evaluation due to invasive nature of obtaining a specimen.”
- Rhinomanometry: “Rhinomanometry is useful in distinguishing between structural and soft tissue causes of obstruction, when history and examination findings are not congruent, as well as a research tool. Better with individual nasal cavity assessment and four-phase rhinomanometry.”
- Acoustic rhinometry: “Acoustic rhinometry is most useful in research setting as opposed to as a clinical diagnostic tool.”
- Peak nasal inspiratory flow: “Use in conjunction with patient reported outcome measures to improve utility.”
- Nitric oxide measurements: “There is inconsistent evidence in the ability of FeNO or nNO to differentiate adults and children with AR and non-allergic rhinitis. Most studies were of low evidence or small impact. There is no agreed upon cut-off value when performing FeNO or nNO for the diagnosis of AR... History and physical, diagnostic skin testing, or sIgE testing should be the first-line evaluation of AR. FeNO or nasal NO testing may

provide additional diagnostic information if necessary but should not be routinely employed for AR diagnosis.”

The National Academies of Science, Engineering and Medicine

The National Academies of Science, Engineering and Medicine convened an expert committee to review the science and management practices of food allergy. Overall, they found that:

- “Currently, no simple diagnostic tests exist for food allergy.”
- “Food allergy evaluation procedures include a medical history and physical examination, and also may include food-specific skin prick test, food-specific serum immunoglobulin E test, diagnostic food elimination diet, and oral food challenge (OFC). Selection of the specific tests needs to be individualized based on the medical history of each patient.”
- “The BAT shows promising preliminary data, the potential utility is recognized and will require additional validation and standardization. “Guidelines suggest not using the BAT clinically on the grounds that it is nonstandardized, but recognize its use as a research tool” (NASEM, 2016).

In 2017, the National Academies of Science, Engineering and Medicine convened an expert committee to examine critical issues related to food allergy. Regarding diagnosis and prognosis, the committee notes that “physicians [should] use evidence-based, standardized procedures as the basis for food allergy diagnosis and avoid nonstandardized and unproven procedures.... When food allergy is suspected, the patient should be evaluated by a physician who has the training and experience to select and interpret appropriate diagnostic tests” (Sicherer et al., 2017).

American Academy of Family Physicians (AAFP)

American Academy of Family Physicians recommendations for practice state: “Allergy and immunologic testing can help clarify the diagnosis and guide treatment. Immediate immunoglobulin E (IgE) and delayed T cell–mediated reactions are the main types of allergic responses. The allergens suspected in an immediate IgE-mediated response are identified through serum IgE-specific antibody or skin testing. For patients with an inhalant allergy, skin or IgE-specific antibody testing is preferred. In patients with food allergies, eliminating the suspected allergenic food from the diet is the initial treatment. If this is ineffective, IgE-specific antibody or skin testing can exclude allergens. An oral food challenge should be performed to confirm the diagnosis. Patients with an anaphylactic reaction to an insect sting should undergo IgE-specific antibody or skin testing. Skin testing for penicillin has a high negative predictive value and can help when penicillin administration is indicated and there are limited alternatives. Testing for other drug allergies has less well-determined sensitivity and specificity but can guide the diagnosis. Patch testing can help identify the allergen responsible for contact dermatitis” (Chang & Guarderas, 2018).

European Academy of Allergy and Clinical Immunology (EAACI)

The EAACI published guidelines on “Biomarkers for monitoring the clinical efficacy of allergen Immunotherapy (AIT).” In it, they concluded that “to date, there are no validated and generally accepted candidate biomarkers that are predictive or indicative of the clinical response to AIT.”

However, they did note sIgE/tIgE ratio and IgE-FAB as candidate biomarkers for future research (Shamji et al., 2017).

The EAACI released a position statement on the BAT. In it, they concluded that “Basophil activation test has been established as a routine diagnostic test with standardized allergen preparations in a number of service laboratories... An important next step is the standardization and automation of analysis of BAT. Once that is achieved, it will be possible to do large multicenter trials to characterize the diagnostic performance of BAT and broaden its use as a clinical tool” (Hoffmann et al., 2015).

The EAACI released a Molecular Allergology User's Guide 2.0. While not a formal guideline, the guide provides comprehensive information about molecular allergen testing. In terms of IgE antibody testing, guide notes that “The clinical relevance of allergen-specific IgE detection in a patient’s serum is strictly as a marker for allergic sensitisation (risk for allergy) and it alone cannot predict the probability of an allergic reaction. The determination of the diagnostic sensitivity and specificity of IgE antibody assays will thus remain difficult to definitively determine because of the lack of an absolute (gold standard) method of defining the presence of allergic disease. This means that the clinical relevance of an allergic sensitisation (i.e. presence of allergen-specific IgE) independent of the use of allergen extracts or molecules for diagnostic purposes will ultimately be determined only by the physician and not by the test” (Dramburg et al., 2023).

In terms of basophil activation testing, the guide notes that “The BAT can be useful to confirm the diagnosis of food, venom and respiratory allergies” (Dramburg et al., 2023).

In terms of *In vivo* testing, the guide notes that “provocation tests are especially helpful when discrepancies exist between the clinical history and other *in vivo* or *in vitro* test results, to phenotype patients and to monitor the efficacy of allergen-specific immunotherapy” but further states that “the use of recombinant allergens in provocation tests seems to improve their accuracy; however, it is an unmet need which requires further investigations” (Dramburg et al., 2023).

National Institute for Health and Care Excellence (NICE)

National Institute for Health and Care Excellence published a guideline on asthma, recommending against use of serum total or specific IgE for diagnosing asthma. Specific IgE or prick tests to aeroallergens should be used to identify triggers to asthma after a formal diagnosis has been made (NICE, 2021).

The NICE also released a statement on multiplex allergen testing, particularly “ImmunoCAP ISAC” Although they acknowledge the test’s promise, they state that there is “insufficient evidence to recommend the routine adoption of multiplex allergen testing with ImmunoCAP ISAC 112 to help diagnose allergy and predict the risk of an allergic reaction in people with allergy that is difficult to diagnose, when used with standard clinical assessment” (NICE, 2020).

Regarding the assessment and diagnosis of food allergy in under 19s, NICE published the below recommendations:

For food allergies classified as IgE-mediated:

“Based on the results of the allergy-focused clinical history, if IgE-mediated allergy is suspected, offer the child or young person a skin prick test and/or blood tests for specific IgE antibodies to the suspected foods and likely co-allergens.”

“Tests should only be undertaken by healthcare professionals with the appropriate competencies to select, perform and interpret them.”

“Skin prick tests should only be undertaken where there are facilities to deal with an anaphylactic reaction.”

“Choose between a skin prick test and a specific IgE antibody blood test based on:

- the results of the allergy-focused clinical history **and**
- whether the test is suitable for, safe for and acceptable to the child or young person (or their parent or carer) **and**
- the available competencies of the healthcare professional to undertake the test and interpret the results.”

“Do not carry out allergy testing without first taking an allergy-focused clinical history. Interpret the results of tests in the context of information from the allergy-focused clinical history.”

“Do not use atopy patch testing or oral food challenges to diagnose IgE-mediated food allergy in primary care or community settings” (NICE, 2011).

VII. Applicable State and Federal Regulations

DISCLAIMER: If there is a conflict between this Policy and any relevant, applicable government policy for a particular member [e.g., Local Coverage Determinations (LCDs) or National Coverage Determinations (NCDs) for Medicare and/or state coverage for Medicaid], then the government policy will be used to make the determination. For the most up-to-date Medicare policies and coverage, please visit the Medicare search website: <http://www.cms.gov/medicare-coverage-database/search.aspx>. For the most up-to-date Medicaid policies and coverage, visit the applicable state Medicaid website.

Food and Drug Administration (FDA)

Many labs have developed specific tests that they must validate and perform in house. These laboratory-developed tests (LDTs) are regulated by the Centers for Medicare and Medicaid (CMS) as high-complexity tests under the Clinical Laboratory Improvement Amendments of 1988 (CLIA '88). LDTs are not approved or cleared by the U. S. Food and Drug Administration; however, FDA clearance or approval is not currently required for clinical use.

VIII. Applicable CPT/HCPCS Procedure Codes

CPT	Code Description
82784	Gammaglobulin (immunoglobulin); IgA, IgD, IgG, IgM, each
82785	Gammaglobulin (immunoglobulin); IgE
82787	Gammaglobulin (immunoglobulin); immunoglobulin subclasses (eg, IgG1, 2, 3, or 4), each
83516	Immunoassay for analyte other than infectious agent antibody or infectious agent antigen; qualitative or semiquantitative, multiple step method
86001	Allergen specific IgG quantitative or semiquantitative, each allergen

86003	Allergen specific IgE; quantitative or semiquantitative, crude allergen extract, each
86005	Allergen specific IgE; qualitative, multiallergen screen (eg, disk, sponge, card)
86008	Allergen specific IgE; quantitative or semiquantitative, recombinant or purified component, each
88184	Flow cytometry, cell surface, cytoplasmic, or nuclear marker, technical component only; first marker
88185	Flow cytometry, cell surface, cytoplasmic, or nuclear marker, technical component only; each additional marker
0165U	Peanut allergen-specific quantitative assessment of multiple epitopes using enzyme-linked immunosorbent assay (ELISA), blood, individual epitope results and probability of peanut allergy Proprietary test: VeriMAP™ Peanut Dx – Bead-based Epitope Assay Lab/Manufacturer: AllerGenis™ Clinical Laboratory
0178U	Peanut allergen-specific quantitative assessment of multiple epitopes using enzyme-linked immunosorbent assay (ELISA), blood, report of minimum eliciting exposure for a clinical reaction Proprietary test: VeriMAP™ Peanut Sensitivity - Bead Based Epitope Assay Lab/Manufacturer: AllerGenis™ Clinical Laboratory

Current Procedural Terminology© American Medical Association. All Rights reserved.

Procedure codes appearing in Medical Policy documents are included only as a general reference tool for each policy. They may not be all-inclusive.

IX. Evidence-based Scientific References

- AAAAI. (2012). Some Tests and Procedures Are Over or Misused to Diagnose and Treat Allergies, Asthma and Immunologic Disorders.
<https://www.aaaai.org/Aaaai/media/MediaLibrary/PDF%20Documents/Media/Choosing-WiselyT.pdf>
- AAP. (2012). Allergy Testing in Childhood: Using Allergen-Specific IgE Tests.
<https://pdfs.semanticscholar.org/51a4/dfa0a84e4dc8b0893529a6197bd9b94bdbfe.pdf>
- Abrams, E. M., Chan, E. S., & Sicherer, S. (2020). Peanut Allergy: New Advances and Ongoing Controversies. *Pediatrics*, 145(5), e20192102. <https://doi.org/10.1542/peds.2019-2102>
- Adams, P. F., Kirzinger, W. K., & Martinez, M. (2013). Summary health statistics for the U.S. population: National Health Interview Survey, 2012. *Vital Health Stat 10(259)*, 1-95.
<https://pubmed.ncbi.nlm.nih.gov/25116371/>
- Ansotegui, Melioli, G., Canonica, G. W., Caraballo, L., Villa, E., Ebisawa, M., Passalacqua, G., Savi, E., Ebo, D., Gómez, R. M., Luengo Sánchez, O., Oppenheimer, J. J., Jensen-Jarolim, E., Fischer, D. A., Haahtela, T., Antila, M., Bousquet, J. J., Cardona, V., Chiang, W. C., . . . Zuberbier, T. (2020). IgE allergy diagnostics and other relevant tests in allergy, a World Allergy Organization position paper. *World Allergy Organ J*, 13(2), 100080.
<https://doi.org/10.1016/j.waojou.2019.100080>
- Ansotegui, Melioli, G., Canonica, G. W., Gomez, R. M., Jensen-Jarolim, E., Ebisawa, M., Luengo, O., Caraballo, L., Passalacqua, G., Poulsen, L., Savi, E., Zuberbier, T., Villa, E., & Oppenheimer, J. (2020). A WAO - ARIA - GA(2)LEN consensus document on molecular-

- based allergy diagnosis (PAMD@): Update 2020. *World Allergy Organ J*, 13(2), 100091. <https://doi.org/10.1016/j.waojou.2019.100091>
- Bahri, R., Custovic, A., Korosec, P., Tsoumani, M., Barron, M., Wu, J., Sayers, R., Weimann, A., Ruiz-Garcia, M., Patel, N., Robb, A., Shamji, M. H., Fontanella, S., Silar, M., Mills, E., Simpson, A., Turner, P. J., & Bulfone-Paus, S. (2018). Mast cell activation test in the diagnosis of allergic disease and anaphylaxis. In *J Allergy Clin Immunol* (Vol. 142, pp. 485-496 e416). <https://doi.org/10.1016/j.jaci.2018.01.043>
- Bernstein, I. L., Li, J. T., Bernstein, D. I., Hamilton, R., Spector, S. L., Tan, R., Sicherer, S., Golden, D. B., Khan, D. A., Nicklas, R. A., Portnoy, J. M., Blessing-Moore, J., Cox, L., Lang, D. M., Oppenheimer, J., Randolph, C. C., Schuller, D. E., Tilles, S. A., Wallace, D. V., . . . Weber, R. (2008). Allergy diagnostic testing: an updated practice parameter. *Ann Allergy Asthma Immunol*, 100(3 Suppl 3), S1-148. [https://doi.org/10.1016/s1081-1206\(10\)60305-5](https://doi.org/10.1016/s1081-1206(10)60305-5)
- Beyer, K., & Teuber, S. S. (2005). Food allergy diagnostics: scientific and unproven procedures. *Curr Opin Allergy Clin Immunol*, 5(3), 261-266. <https://doi.org/10.1097/01.all.0000168792.27948.f9>
- BioSpace. (2021). *AllerGenis Allergy Diagnostic Company to Present Significant Milestones at the Biotech Showcase(TM) 2021*. <https://www.biospace.com/article/allergen-is-allergy-diagnostic-company-to-present-significant-milestones-at-the-biotech-showcase-tm-2021/>
- Boyce, J. A., Assa'ad, A., Burks, A. W., Jones, S. M., Sampson, H. A., Wood, R. A., Plaut, M., Cooper, S. F., Fenton, M. J., Arshad, S. H., Bahna, S. L., Beck, L. A., Byrd-Bredbenner, C., Camargo, C. A., Jr., Eichenfield, L., Furuta, G. T., Hanifin, J. M., Jones, C., Kraft, M., . . . Schwaninger, J. M. (2010). Guidelines for the Diagnosis and Management of Food Allergy in the United States: Summary of the NIAID-Sponsored Expert Panel Report. *J Allergy Clin Immunol*, 126(6), 1105-1118. <https://doi.org/10.1016/j.jaci.2010.10.008>
- Caglayan Sozmen, S., Povesi Dascola, C., Gioia, E., Mastrorilli, C., Rizzuti, L., & Caffarelli, C. (2015). Diagnostic accuracy of patch test in children with food allergy. *Pediatr Allergy Immunol*, 26(5), 416-422. <https://doi.org/10.1111/pai.12377>
- Cardona, V., Ansotegui, I. J., Ebisawa, M., El-Gamal, Y., Fernandez Rivas, M., Fineman, S., Geller, M., Gonzalez-Estrada, A., Greenberger, P. A., Sanchez Borges, M., Senna, G., Sheikh, A., Tanno, L. K., Thong, B. Y., Turner, P. J., & Worm, M. (2020). World allergy organization anaphylaxis guidance 2020. *World Allergy Organ J*, 13(10), 100472. <https://doi.org/10.1016/j.waojou.2020.100472>
- Carlsson, M., Thorell, L., Sjolander, A., & Larsson-Faria, S. (2015). Variability of total and free IgE levels and IgE receptor expression in allergic subjects in and out of pollen season. *Scand J Immunol*, 81(4), 240-248. <https://doi.org/10.1111/sji.12270>
- Cell Science Systems. (2023). Identify food and chemical sensitivities with the Alcat Test. <https://cellsciencesystems.com/providers/alcat-test/>
- Chang, & Guarderas. (2018). Allergy Testing: Common Questions and Answers. *Am Fam Physician*, 98(1), 34-39. <https://www.aafp.org/pubs/afp/issues/2018/0701/p34.html>
- Chow, A. W., Benninger, M. S., Brook, I., Brozek, J. L., Goldstein, E. J., Hicks, L. A., Pankey, G. A., Seleznick, M., Volturo, G., Wald, E. R., & File, T. M., Jr. (2012). IDSA clinical practice guideline for acute bacterial rhinosinusitis in children and adults. *Clin Infect Dis*, 54(8), e72-e112. <https://doi.org/10.1093/cid/cir1043>
- Davila, I., Valero, A., Entrenas, L. M., Valveny, N., & Herraes, L. (2015). Relationship between serum total IgE and disease severity in patients with allergic asthma in Spain. *J Investig Allergol Clin Immunol*, 25(2), 120-127. <https://pubmed.ncbi.nlm.nih.gov/25997305/>

- Depince-Berger, A. E., Sidi-Yahya, K., Jeraiby, M., & Lambert, C. (2017). Basophil activation test: Implementation and standardization between systems and between instruments. *Cytometry A*, 91(3), 261-269. <https://doi.org/10.1002/cyto.a.23078>
- Dramburg, S., Hilger, C., Santos, A. F., de Las Vecillas, L., Aalberse, R. C., Acevedo, N., Aglas, L., Altmann, F., Arruda, K. L., Asero, R., Ballmer-Weber, B., Barber, D., Beyer, K., Biedermann, T., Bilo, M. B., Blank, S., Bosshard, P. P., Breiteneder, H., Brough, H. A., . . . Hoffmann-Sommergruber, K. (2023). EAACI Molecular Allergology User's Guide 2.0. *Pediatr Allergy Immunol*, 34 Suppl 28, e13854. <https://doi.org/10.1111/pai.13854>
- Dykewicz, M. S., Wallace, D. V., Amrol, D. J., Baroody, F. M., Bernstein, J. A., Craig, T. J., Dinakar, C., Ellis, A. K., Finegold, I., Golden, D. B. K., Greenhawt, M. J., Hagan, J. B., Horner, C. C., Khan, D. A., Lang, D. M., Larenas-Linnemann, D. E. S., Lieberman, J. A., Meltzer, E. O., Oppenheimer, J. J., . . . Steven, G. C. (2020). Rhinitis 2020: A practice parameter update. *J Allergy Clin Immunol*, 146(4), 721-767. <https://doi.org/10.1016/j.jaci.2020.07.007>
- FDA. (2016). *Xolair Label*. https://www.accessdata.fda.gov/drugsatfda_docs/label/2016/103976s52251bl.pdf
- Fonacier, L., Bernstein, D. I., Pacheco, K., Holness, D. L., Blessing-Moore, J., Khan, D., Lang, D., Nicklas, R., Oppenheimer, J., Portnoy, J., Randolph, C., Schuller, D., Spector, S., Tilles, S., & Wallace, D. (2015). Contact dermatitis: a practice parameter-update 2015. *J Allergy Clin Immunol Pract*, 3(3 Suppl), S1-39. <https://doi.org/10.1016/j.jaip.2015.02.009>
- Genova. (2023). Allergix® IgG4 Food Antibodies 90 - Serum. <https://www.gdx.net/product/allergix-igg4-food-antibodies-90-food-sensitivity-test-serum>
- Greenhawt, M., Shaker, M., Wang, J., Oppenheimer, J. J., Sicherer, S., Keet, C., Swaggart, K., Rank, M., Portnoy, J. M., Bernstein, J., Chu, D. K., Dinakar, C., Golden, D., Horner, C., Lang, D. M., Lang, E. S., Khan, D. A., Lieberman, J., Stukus, D., & Wallace, D. (2020). Peanut allergy diagnosis: A 2020 practice parameter update, systematic review, and GRADE analysis. *J Allergy Clin Immunol*, 146(6), 1302-1334. <https://doi.org/10.1016/j.jaci.2020.07.031>
- Greer, F. R., Sicherer, S. H., & Burks, A. W. (2019). The Effects of Early Nutritional Interventions on the Development of Atopic Disease in Infants and Children: The Role of Maternal Dietary Restriction, Breastfeeding, Hydrolyzed Formulas, and Timing of Introduction of Allergenic Complementary Foods. *Pediatrics*, 143(4). <https://doi.org/10.1542/peds.2019-0281>
- Hamilton, R. (2023). *Allergen sampling in the environment - UpToDate* (P. S. Creticos & A. Feldweg, Eds.) <https://www.uptodate.com/contents/allergen-sampling-in-the-environment>
- Hamilton, R. G., Matsson, P. N., Hovanec-Burns, D. L., Van Cleve, M., Chan, S., Kober, A., Kleine-Tebbe, J. R., Renz, H., Magnusson, C., & Quicho, R. (2015). Analytical Performance Characteristics, Quality Assurance and Clinical Utility of Immunological Assays for Human IgE Antibodies of Defined Allergen Specificities.(CLSI-ILA20-A3). *Journal of Allergy and Clinical Immunology*, 135(2), AB8. <https://doi.org/10.1016/j.jaci.2014.12.961>
- Hammond, C., & Lieberman, J. A. (2018). Unproven Diagnostic Tests for Food Allergy. *Immunol Allergy Clin North Am*, 38(1), 153-163. <https://doi.org/10.1016/j.iac.2017.09.011>
- He, Y. T., & Reisacher, W. R. (2019). Sensitivity, specificity, and predictive value of oral mucosal brush biopsy for the diagnosis of peanut allergy. *Int Forum Allergy Rhinol*, 9(6), 624-628. <https://doi.org/10.1002/alr.22302>

- Hemmings, O., Kwok, M., McKendry, R., & Santos, A. F. (2018). Basophil Activation Test: Old and New Applications in Allergy. *Current Allergy and Asthma Reports, 18*(12), 77. <https://doi.org/10.1007/s11882-018-0831-5>
- Hoffmann, H. J., Santos, A. F., Mayorga, C., Nopp, A., Eberlein, B., Ferrer, M., Rouzaire, P., Ebo, D. G., Sabato, V., Sanz, M. L., Pecaric-Petkovic, T., Patil, S. U., Hausmann, O. V., Shreffler, W. G., Korosec, P., & Knol, E. F. (2015). The clinical utility of basophil activation testing in diagnosis and monitoring of allergic disease. *Allergy, 70*(11), 1393-1405. <https://doi.org/10.1111/all.12698>
- Jackson, K. D., Howie, L. D., Akinbami, L. J., & CDC. (2013). Trends in Allergic Conditions Among Children: United States, 1997-2011. NCHS Data Brief. No 121. <https://www.cdc.gov/nchs/products/databriefs/db121.htm>
- Kim, S. Y., Kim, J. H., Jang, Y. S., Choi, J. H., Park, S., Hwang, Y. I., Jang, S. H., & Jung, K. S. (2016). The Basophil Activation Test Is Safe and Useful for Confirming Drug-Induced Anaphylaxis. *Allergy Asthma Immunol Res, 8*(6), 541-544. <https://doi.org/10.4168/aair.2016.8.6.541>
- Klemans, R. J., van Os-Medendorp, H., Blankestijn, M., Bruijnzeel-Koomen, C. A., Knol, E. F., & Knulst, A. C. (2015). Diagnostic accuracy of specific IgE to components in diagnosing peanut allergy: a systematic review. *Clin Exp Allergy, 45*(4), 720-730. <https://doi.org/10.1111/cea.12412>
- Knight, V., Wolf, M. L., Trikha, A., Curran-Everett, D., Hiserote, M., & Harbeck, R. J. (2018). A comparison of specific IgE and skin prick test results to common environmental allergens using the HYTEC™ 288. *Journal of Immunological Methods, 462*, 9-12. <https://doi.org/10.1016/j.jim.2018.07.005>
- Kowal, K., & DuBuske, L. (2021, 05/03/2021). *Overview of in vitro allergy tests*. <https://www.uptodate.com/contents/overview-of-in-vitro-allergy-tests>
- Kowal, K., & DuBuske, L. (2022, 11/30/2022). *Overview of skin testing for allergic disease - UpToDate*. [https://www.uptodate.com/contents/overview-of-skin-testing-for-allergic-disease - UpToDate](https://www.uptodate.com/contents/overview-of-skin-testing-for-allergic-disease-UpToDate). [https://www.uptodate.com/contents/overview-of-skin-testing-for-allergic-disease - UpToDate](https://www.uptodate.com/contents/overview-of-skin-testing-for-allergic-disease-UpToDate)
- Lieberman, P., Nicklas, R. A., Randolph, C., Oppenheimer, J., Bernstein, D., Bernstein, J., Ellis, A., Golden, D. B., Greenberger, P., Kemp, S., Khan, D., Ledford, D., Lieberman, J., Metcalfe, D., Nowak-Wegrzyn, A., Sicherer, S., Wallace, D., Blessing-Moore, J., Lang, D., . . . Tilles, S. A. (2015). Anaphylaxis--a practice parameter update 2015. *Ann Allergy Asthma Immunol, 115*(5), 341-384. <https://doi.org/10.1016/j.anai.2015.07.019>
- Mowad, C. M. (2006). Patch testing: pitfalls and performance. *Curr Opin Allergy Clin Immunol, 6*(5), 340-344. <https://doi.org/10.1097/01.all.0000244794.03239.8e>
- NASEM. (2016). *Finding a Path to Safety in Food Allergy: Assessment of the Global Burden, Causes, Prevention, Management, and Public Policy* (Finding a Path to Safety in Food Allergy: Assessment of the Global Burden, Causes, Prevention, Management, and Public Policy, Issue. <http://dx.doi.org/10.17226/23658>
- Nelson, H. S. (2001). Variables in Allergy Skin Testing. *Immunology and Allergy Clinics, 21*(2), 281-290. [https://doi.org/10.1016/S0889-8561\(05\)70206-X](https://doi.org/10.1016/S0889-8561(05)70206-X)
- NICE. (2011, February 23, 2011). *Food allergy in under 19s: assessment and diagnosis*. National Institute for Health and Care Excellence (NICE). <https://www.nice.org.uk/guidance/cg116/chapter/Recommendations#ige-mediated-food-allergy>
- NICE. (2020). ImmunoCAP ISAC 112 for multiplex allergen testing. <https://www.nice.org.uk/guidance/dg24/chapter/1-Recommendations>

- NICE. (2021). Asthma: diagnosis, monitoring and chronic asthma management. <https://www.nice.org.uk/guidance/ng80>
- Oppenheimer, J., & Nelson, H. S. (2006). Skin testing: a survey of allergists. *Ann Allergy Asthma Immunol*, 96(1), 19-23. [https://doi.org/10.1016/s1081-1206\(10\)61034-4](https://doi.org/10.1016/s1081-1206(10)61034-4)
- Pawankar, R., Holgate, S. T., Canonica, G. W., Lockey, R. F., & Blaiss, M. S. (2013). *WAO White Book on Allergy | World Allergy Organization*. <https://www.forskasverige.se/wp-content/uploads/WAO-WhiteBook-2013.pdf>
- Rietschel, R. L. (1997). COMPARISON OF ALLERGIC AND IRRITANT CONTACT DERMATITIS. *Immunology and Allergy Clinics*, 17(3), 359-364. [https://doi.org/10.1016/S0889-8561\(05\)70314-3](https://doi.org/10.1016/S0889-8561(05)70314-3)
- Sampson, H. A., Aceves, S., Bock, S. A., James, J., Jones, S., Lang, D., Nadeau, K., Nowak-Wegrzyn, A., Oppenheimer, J., Perry, T. T., Randolph, C., Sicherer, S. H., Simon, R. A., Vickery, B. P., Wood, R., Bernstein, D., Blessing-Moore, J., Khan, D., Nicklas, R., . . . Wallace, D. (2014). Food allergy: a practice parameter update-2014. *J Allergy Clin Immunol*, 134(5), 1016-1025.e1043. <https://doi.org/10.1016/j.jaci.2014.05.013>
- Santos, A. F., Douiri, A., Becares, N., Wu, S. Y., Stephens, A., Radulovic, S., Chan, S. M., Fox, A. T., Du Toit, G., Turcanu, V., & Lack, G. (2014). Basophil activation test discriminates between allergy and tolerance in peanut-sensitized children. *J Allergy Clin Immunol*, 134(3), 645-652. <https://doi.org/10.1016/j.jaci.2014.04.039>
- Santos, A. F., Du Toit, G., Douiri, A., Radulovic, S., Stephens, A., Turcanu, V., & Lack, G. (2015). Distinct parameters of the basophil activation test reflect the severity and threshold of allergic reactions to peanut. *J Allergy Clin Immunol*, 135(1), 179-186. <https://doi.org/10.1016/j.jaci.2014.09.001>
- Santos, A. F., & Lack, G. (2016). Basophil activation test: food challenge in a test tube or specialist research tool? *Clin Transl Allergy*, 6, 10. <https://doi.org/10.1186/s13601-016-0098-7>
- Santos, A. F., & Shreffler, W. G. (2017). Road map for the clinical application of the basophil activation test in food allergy. *Clin Exp Allergy*, 47(9), 1115-1124. <https://doi.org/10.1111/cea.12964>
- Shamji, M. H., Kappen, J. H., Akdis, M., Jensen-Jarolim, E., Knol, E. F., Kleine-Tebbe, J., Bohle, B., Chaker, A. M., Till, S. J., Valenta, R., Poulsen, L. K., Calderon, M. A., Demoly, P., Pfaar, O., Jacobsen, L., Durham, S. R., & Schmidt-Weber, C. B. (2017). Biomarkers for monitoring clinical efficacy of allergen immunotherapy for allergic rhinoconjunctivitis and allergic asthma: an EAACI Position Paper. *Allergy*, 72(8), 1156-1173. <https://doi.org/10.1111/all.13138>
- Sicherer, S. (2017). *New guidelines detail use of 'infant-safe' peanut to prevent allergy*. <https://www.aappublications.org/news/2017/01/05/PeanutAllergy010517>
- Sicherer, S. H., Allen, K., Lack, G., Taylor, S. L., Donovan, S. M., & Oria, M. (2017). Critical Issues in Food Allergy: A National Academies Consensus Report. *Pediatrics*, 140(2). <https://doi.org/10.1542/peds.2017-0194>
- Siroux, V., Lupinek, C., Resch, Y., Curin, M., Just, J., Keil, T., Kiss, R., Lodrup Carlsen, K., Melen, E., Nadif, R., Pin, I., Skrindo, I., Vrtala, S., Wickman, M., Anto, J. M., Valenta, R., & Bousquet, J. (2017). Specific IgE and IgG measured by the MeDALL allergen-chip depend on allergen and route of exposure: The EGEA study. *J Allergy Clin Immunol*, 139(2), 643-654.e646. <https://doi.org/10.1016/j.jaci.2016.05.023>

- Soares-Weiser, K., Takwoingi, Y., Panesar, S. S., Muraro, A., Werfel, T., Hoffmann-Sommergruber, K., Roberts, G., Halken, S., Poulsen, L., van Ree, R., Vlieg-Boerstra, B. J., & Sheikh, A. (2014). The diagnosis of food allergy: a systematic review and meta-analysis. *Allergy*, *69*(1), 76-86. <https://doi.org/10.1111/all.12333>
- Sookrung, N., Jotikaprasardhna, P., Bunnag, C., Chaicumpa, W., & Tungtrongchitr, A. (2019). Concordance of skin prick test and serum-specific IgE to locally produced component-resolved diagnostics for cockroach allergy. *Ann Allergy Asthma Immunol*, *122*(1), 93-98. <https://doi.org/10.1016/j.anai.2018.09.463>
- Spiriplex. (2023). Allergies. <https://spiriplex.com/allergies/>
- Stokes, J., & Casale, T. (2022, 11/22/2022). *The relationship between IgE and allergic disease*. <https://www.uptodate.com/contents/the-relationship-between-ige-and-allergic-disease>
- Suárez-Fariñas, M., Suprun, M., Kearney, P., Getts, R., Grishina, G., Hayward, C., Luta, D., Porter, A., Witmer, M., du Toit, G., Lack, G., Chinthrajah, R. S., Galli, S. J., Nadeau, K., & Sampson, H. A. (2021). Accurate and reproducible diagnosis of peanut allergy using epitope mapping. *Allergy*, *n/a*(n/a). <https://doi.org/10.1111/all.14905>
- Tannert, L. K., Mortz, C. G., Skov, P. S., & Bindslev-Jensen, C. (2017). Positive Skin Test or Specific IgE to Penicillin Does Not Reliably Predict Penicillin Allergy. *J Allergy Clin Immunol Pract*, *5*(3), 676-683. <https://doi.org/10.1016/j.jaip.2017.03.014>
- Togias, A., Cooper, S. F., Acebal, M. L., Assa'ad, A., Baker, J. R., Jr., Beck, L. A., Block, J., Byrd-Bredbenner, C., Chan, E. S., Eichenfield, L. F., Fleischer, D. M., Fuchs, G. J., 3rd, Furuta, G. T., Greenhawt, M. J., Gupta, R. S., Habich, M., Jones, S. M., Keaton, K., Muraro, A., . . . Boyce, J. A. (2017). Addendum guidelines for the prevention of peanut allergy in the United States: Report of the National Institute of Allergy and Infectious Diseases-sponsored expert panel. *J Allergy Clin Immunol*, *139*(1), 29-44. <https://doi.org/10.1016/j.jaci.2016.10.010>
- Tschopp, J. M., Sistek, D., Schindler, C., Leuenberger, P., Perruchoud, A. P., Wuthrich, B., Brutsche, M., Zellweger, J. P., Karrer, W., & Brandli, O. (1998). Current allergic asthma and rhinitis: diagnostic efficiency of three commonly used atopic markers (IgE, skin prick tests, and Phadiatop). Results from 8329 randomized adults from the SAPALDIA Study. Swiss Study on Air Pollution and Lung Diseases in Adults. *Allergy*, *53*(6), 608-613. <https://doi.org/10.1111/j.1398-9995.1998.tb03937.x>
- Usmani, N., & Wilkinson, S. M. (2007). Allergic skin disease: investigation of both immediate- and delayed-type hypersensitivity is essential. *Clin Exp Allergy*, *37*(10), 1541-1546. <https://doi.org/10.1111/j.1365-2222.2007.02805.x>
- Werther, R. L., Choo, S., Lee, K. J., Poole, D., Allen, K. J., & Tang, M. L. (2012). Variability in Skin Prick Test Results Performed by Multiple Operators Depends on the Device Used. *World Allergy Organ J*, *5*(12), 200-204. <https://doi.org/10.1097/WOX.0b013e31827e6513>
- Wise, S. K., Damask, C., Roland, L. T., Ebert, C., Levy, J. M., Lin, S., Luong, A., Rodriguez, K., Sedaghat, A. R., Toskala, E., Villwock, J., Abdullah, B., Akdis, C., Alt, J. A., Ansotegui, I. J., Azar, A., Baroody, F., Benninger, M. S., Bernstein, J., . . . Zhang, L. (2023). International consensus statement on allergy and rhinology: Allergic rhinitis - 2023. *Int Forum Allergy Rhinol*, *13*(4), 293-859. <https://doi.org/10.1002/alr.23090>
- Wuthrich, B. (2005). Unproven techniques in allergy diagnosis. *J Invest Allergol Clin Immunol*, *15*(2), 86-90. <https://pubmed.ncbi.nlm.nih.gov/16047707/>
- Zug, K. A., Pham, A. K., Belsito, D. V., DeKoven, J. G., DeLeo, V. A., Fowler, J. F., Jr., Fransway, A. F., Maibach, H. I., Marks, J. G., Jr., Mathias, C. G., Pratt, M. D., Sasseville, D.,

Storrs, F. J., Taylor, J. S., Warshaw, E. M., & Zirwas, M. J. (2014). Patch testing in children from 2005 to 2012: results from the North American contact dermatitis group. *Dermatitis*, 25(6), 345-355. <https://doi.org/10.1097/der.0000000000000083>

X. Review/Revision History

Effective Date	Summary
08/01/2024	Reviewed and Updated: Updated the background, guidelines and recommendations, and evidence-based scientific references. Literature review did not necessitate any modifications to coverage criteria. The following edits were made for clarity and consistency:CC3 edited for clarity and consistency to allow enhanced enforcement.CC7 and CC8 were wrapped into a single DNMCC criterion for individuals with signs and symptoms of allergies. Now reads: “7) For individuals with signs or symptoms of allergies, basophil activation flow cytometry testing (BAT) and in-vitro testing of IgG, IgA, IgM, and/or IgD DO NOT MEET COVERAGE CRITERIA.”
07/01/2023	Annual Review: Literature review did not necessitate changes to coverage criteria. Policy edited for clarity and consistency.
07/01/2022	<p>Off-Cycle Review: Literature review necessitated the addition of a new CC4 that reads "In individuals with positive food allergen results, annual re-testing for the same food allergens MEETS COVERAGE CRITERIA in children and adolescents to monitor for allergy resolution.”</p> <p>Edited CC5 for clarity with the addition of new CC “Except as specified in criteria 4 (above), routine re-testing for allergies to the same allergens DOES NOT MEET COVERAGE CRITERIA in the absence of a new clinical presentation.”</p> <p>Removed “When in-vitro testing is ordered, the medical record must clearly document the indication and why it is being used instead of skin testing.” from CC1a. Now reads: “a) In lieu of skin testing for an INITIAL allergy screen.” Reference source “reevaluation is usually performed yearly.”</p> <p>CPT Changes: Removed code 83520</p>
11/04/2021	Initial Policy Implementation